

FR_TERBIT.pdf

by

Submission date: 24-Apr-2023 11:16AM (UTC+0700)

Submission ID: 2073587230

File name: FR_TERBIT.pdf (555.98K)

Word count: 6408

Character count: 30691

43
Rice bran oil extraction by ethanol: optimization of γ -oryzanol and polyphenol

1,* Mas'ud, F., 1 Banggalino, H., 1 Yusuf, M., 2 Suhardi, S. and 3 Sayuti, M.

1 Department of Chemical Engineering, Politeknik Negeri Ujung Pandang, Makassar, South Sulawesi, 90245, Indonesia

2 Department of Agricultural Engineering, Hasanuddin University, Makassar, South Sulawesi, 90245, Indonesia

3 Department of Animal Husbandry, Agricultural Faculty, Gorontalo State University, Gorontalo, 96128, Indonesia

Article history:

Received: 8 January 2022

Received in revised form: 25 February 2022

Accepted: 19 April 2022

Available Online: 22 April 2023

Keywords:

Rice bran oil,
Reflux method,
Ethanol,
Response surface
methodology

17

DOI:

[https://doi.org/10.26656/fr.2017.7\(2\).011](https://doi.org/10.26656/fr.2017.7(2).011)

Abstract

Rice bran oil (RBO) contains γ -oryzanol and polyphenols which are very beneficial for health. The work investigated the effect of temperature ($^{\circ}\text{C}$) (X_1), time (hr) (X_2), and volume of ethanol (mL) (X_3) as a solvent on the oil yield (%) (Y_1), γ -oryzanol (Y_2), and total polyphenol content (TPC) of RBO (Y_3) with apply reflux extraction method. The study was conducted using Response Surface Methodology (RSM) based on Central Composite Design (CCD) model with three levels and six centre points. The data were analyzed using Design-Expert 10 software to create and evaluate models and to plot the response curves of the 3-dimensional surfaces. The maximum of $Y_1 = 10.4\%$, $Y_2 = 195.49 \text{ mg.L}^{-1}$, and $Y_3 = 1.02 \text{ mg gallic acid equivalent (GAE).g}^{-1}$ were achieved under the optimum conditions process of X_1 , X_2 , and X_3 were 78.7°C , 1.6 hrs, and 496.5 mL ethanol/100 g rice bran, respectively. The average of Y_1 , Y_2 and Y_3 on verified models were $16.1 \pm 0.66\%$, $189.12 \pm 1.18 \text{ mg.L}^{-1}$, and $0.91 \pm 0.54 \text{ mg GAE.g}^{-1}$, respectively. These indicated that this method is very promising to be applied in industry and as an effort to apply the concept of "green" solvents that is safe for humans and the environment.

1. Introduction

Rice bran, a low-value co-product obtained from rice processing, represents a potential source of healthy products due to its nutritional aspects, especially the oil it contains. RBO is rich in natural antioxidants or various bioactive compounds, including γ -oryzanol, phytosterols, sterol esters, triterpene alcohols, tocopherols, tocotrienols and other phenolic compounds (Patel and Naik, 2004). These minor compounds have been cited in the scientific literature as powerful antioxidant agents that are effective in preventing degenerative diseases (Lerma-García *et al.*, 2009). The antioxidants of RBO have the potential use as additives to improve the storage stability of foods (Iqbal *et al.*, 2005).

Several studies have reported the potential of RBO. Therefore, this study was directed at optimizing the extraction process. In recent years, the oil industry has shown increased interest in alternative solvents because of environmental and safety concerns. Ethanol has gained attention as a potential solvent for vegetable oils (Rodrigues *et al.*, 2010). As a result of study before,

ethanol was feasible to use for RBO extraction by maceration method (Mas'ud *et al.*, 2019), furthermore, Mas'ud *et al.* (2020) reported that ethanol has proven as an effective and efficient solvent for extracting oil with high yield, and it feasible to be applied on the extraction of mango seed kernel oil (Mas'ud *et al.*, 2021). Ethanol has been widely applied as a viable solvent due to its ease of recovery and low cost in an application and is classified as an environmentally friendly green solvent (Péres *et al.*, 2006). Green solvents have several benefits such as biodegradability, low toxicity, non-flammability, and renewability, making them potential candidates in separation/extraction science (Mas'ud *et al.*, 2017).

Do *et al.* (2014) report that ethanol is a good solvent for polyphenols, and in this work, the feasibility of ethanol as a solvent for RBO on reflux method was investigated from the perspective of the extracted oil, γ -oryzanol, and polyphenols. Previous studies showed that temperature, time, and solvent were the main parameters affecting the extraction yield from different sources (Minjares-Fuentes *et al.*, 2014). These parameters also

*Corresponding author.

Email: fajri888@polupg.ac.id

had a significant influence on the endogenous bioactive compounds, such as certain from the TPC (Rodrigues *et al.*, 2008; Chanioti *et al.*, 2016). Oliveira *et al.* (2012) have used ethanol in RBO extraction to study its effect on γ -oryzanol and tocopherols. It was observed that γ -oryzanol and tocopherols behave in different ways during the extraction process, and ethanol is feasible to obtain enriched oil when this renewable solvent is used.

Based on the advantages of ethanol as solvent, this study is directed to investigate the effect of temperature, time, and volume of ethanol on the oil yield, γ -oryzanol, and total polyphenol content (TPC) of RBO, and apply the reflux extraction method using RSM based CCD model to optimize the RBO extraction process, standardize, and analyses the models.

2. Materials and methods

2.1 Material

Celebes rice, local rice of Indonesia, as samples were obtained from milling rice grain in a local grinding mill in Makassar, Indonesia from March to April 2021. Ethanol (wt.%) was purchased from a local chemical shop, γ -oryzanol standard from Sigma-Aldrich Co. All chemicals were from Merck, Germany.

2.2 Preparation of rice bran and extraction oil

Freshly milled rice bran was directly collected from the milling system in polyethylene bags, the rice bran was screened through a 60-mesh sieve to have a uniform particle size and stabilized at autoclave (Hiclave HV-85 Hirayama) at 100°C for 15 mins for inactivating endogenous lipase. For extraction oil, each experimental unit weighed 100 g of rice bran in the reactor 1.0 L four neck flasks, RBO was extracted by reflux method using a heating mantle connected with the thermometer set, agitator on the top with the speed of 100 rpm, the residue was separated by centrifugation (refrigerated AX-521 centrifuge) at a speed of 3500 rpm for 20 mins. The liquid part is accommodated in the flask evaporator and ethanol as solvent is removed on a rotary evaporator Buchi R-215 incorporates vacuum Pump V-700. RBO obtained was packaged in a dark glass bottle and stored in a freezer for analysis. The percentage of oil yield was calculated by equation (1) according to Sani (2014):

$$\text{Oil yield (\%)} = \frac{\text{RBO (g)}}{\text{Rice bran sample (g)}} \times 100 \quad (1)$$

2.3 Analysis of γ -oryzanol

Preparation of γ -oryzanol standard: 0.05 g of the pure γ -oryzanol was dissolved in ethyl acetate in a 100 mL flask, and diluted to respectively 100, 200, 300, 400,

and 500 ppm at a 50 mL flask, homogenized by vortex, inserted in the vial GC-MS (gas chromatography-mass spectrometry). Preparation of sample of RBO for γ -oryzanol analysis: 0.012 g RBO dissolved in 2 mL ethyl acetate, homogenized by vortex, and inserted into the vial bottle GC-MS. Quantification of γ -oryzanol of RBO: γ -oryzanol of RBO was performed on a GC-MS QP2010 by Shimadzu equipped with a split/splitless injector. Separations were achieved using a Rxi SH-5Sil MS capillary column (30 m, 0.25 mm ID, 0.25 mm film thickness). Helium was used as the carrier gas at flow rates of 14.0 mL/min and a split ratio of 1:10. The oven temperature was programmed at 110°C for a hold of 2 mins and increased to 200°C at a rate of 10°C/min and hold at the final temperature for 9 mins. LabSolutions software was used to control the operation of GC-MS. MS spectra were obtained at range width m/z 40-450, interface temperature 280°C, and ion source temperature 200°C. γ -oryzanol of RBO peaks was identified by comparing their retention time and equivalent chain length with respect to the standard.

2.4 Determination of total phenolic content

The analysis method of TPC according to Brand-Williams *et al.* (1995), 0.25 mL RBO was mixed with 2 mL of 10% Folin Ciocalteu reagent and 1.6 mL of 7.5% Na_2CO_3 and left at room temperature for 30 mins. The mixing solution was measured absorbance at 750 nm by ultraviolet-visible spectrophotometer (UV 1800, Shimadzu, Japan). Gallic acid was used as a standard and the TPC were expressed as mg gallic acid equivalents (GAE).g⁻¹.

2.5 Experimental design and statistical analysis

The study was conducted using RSM to optimize the temperature (°C) (X_1), time (hr) (X_2), and volume of ethanol (mL) (X_3) to maximize the oil yield (%) (Y_1), γ -oryzanol (mg. L⁻¹) (Y_2), and TPC (mg GAE.g⁻¹) (Y_3) of RBO as responses. The middle values were determined based on the results of the preliminary study, i.e., 80°C, 1.5 hrs, and 450 mL ethanol, respectively. The lower and upper limits for each treatment were 70°C and 90°C for X_1 , 1 hr and 2 hrs for X_2 , 350 mL and 550 mL ethanol for X_3 (Table 1). The CCD consists of 14 experimental units and 6 replications of the centre point. Replication of the centre point was aimed to evaluate the pure error variance as the experimental error and to control the adequacy of the model. To estimate the coefficients of the response function and predict the system's responses, analysis of the experimental results of CCD was realized using empirical second-order polynomial equations as follows Amiri *et al.* (2018).

$$Y = \beta_0 + \sum_{i=1}^k \beta_i X_i + \sum_{i=1}^k \beta_{ii} X_i^2 + \sum_{i=1}^{k-1} \sum_{j=i+1}^k \beta_{ij} X_i X_j + \dots e \quad (2)$$

Where Y is a response, β_0 denotes the constant-coefficient; X_i and X_j the independent factors, β_i , β_{ii} , and β_{ij} the regression coefficients for the linear, quadratic, and interaction effects, respectively; k the number of variables; and e stands for the statistical error occurring to response Y .

Table 1. Five levels of independent variables of the CCD

Independent variables	Star low	Low	Centre	High	Star high
	-1.68	-1	0	1	1.68
X_1 (°C)	63	70	80	90	97
X_2 (h)	0.7	1	1.5	2	2.3
X_3 (mL)	282	350	450	550	619

Statistic software Design-Expert 10 was used to design, analyze, and optimize experimental models. Analysis of variance (ANOVA) was used to validate the statistical significance of the parameters that influence the responses, and the quality of the predicted model (Mohammed *et al.*, 2018). The coefficient of determination (R^2) shows the total predictive performance of the model, it represented the validity and fit quality of the model's quadratic polynomials. A p -value of ≤ 0.05 at a confidence level of 95% and an F -value of the Lack-of-Fit test were used for statistical analysis to evaluate the significance of model statistics (Tan *et al.*, 2017).

3. Results and discussion

3.1 Model fitting

Based on the correlation coefficient value, the quality of the model's responses can be evaluated. According to the ANOVA of Y_1 , Y_2 , and Y_3 , the R^2 values of Y_1 , Y_2 , and Y_3 were 0.94, 0.93, and 0.94, respectively (Table 2). The value of R^2 close to 1 indicates that the models adequately represented the real relationship between the parameters chosen and indicated a reasonable adjustment of the model to experimental data. R^2 is more than 80% implying that the regression model shows good fit (Singh *et al.*, 2018), the model will be significant at a 95% confidence interval if the F test has a p -value of less than 0.05. In the case of lack of fit ($p > F$), the p -value is greater than 0.05 showing the failure of the model in finding data points in the experimental domain. The significance of different terms of each coefficient was determined using the F -value and p -value. According to Yolmeh *et al.* (2014), a

Table 2. Y_1 , Y_2 , and Y_3 of RBO in terms of actual factors

	R^2	a	X_1	X_2	X_3	X_1X_2	X_1X_3	X_2X_3	X_1^2	X_2^2	X_3^2
Y_1 (%)	0.94	-53.87	1.22	9.48	0.05	-0.01	-2.56	-3.68	-6.62	-1.98	-2.03
Y_2 (mg.L ⁻¹)	0.93	-1005.08	17.5	227.72	1.18	0.10	-1.35E-003	-0.17	-0.1	-47.43	-8.20E-004
Y_3 (mg GAE.mg ⁻¹)	0.94	-6.03	0.07	2.40	0.01	-0.01	2.45E-005	-6.78E-004	-2.35E-004	0.29	-9.21

[https://doi.org/10.26656/fr.2017.7\(2\).011](https://doi.org/10.26656/fr.2017.7(2).011)

large F -value and a small p -value would imply a more significant effect on the corresponding response variable. According to Li *et al.* (2014), the model is important and can be used to navigate the design domain.

3.2 Interpretation result of analysis of variance

3.2.1 Oil yield

The result of ANOVA for the response surface quadratic model of Y_1 showed that the F -value model of 17.04 implies the model is significant (Table 3). There is only a 0.01% chance that an F -value this large could occur due to noise. Values of Prob > F of < 0.0001 indicate that the model terms are significant, in this case, the linear term of temperature (X_1), the linear term of time (X_2), the linear term of ethanol volume (X_3), the quadratic terms of temperature (X_1^2), and the quadratic terms of time (X_2^2) are significant model terms. On the other hand, values Prob > F greater than 0.1000 indicate the variable terms are not significant, in this case, the quadratic terms of ethanol volume (X_3^2), the interaction between temperature and time (X_1X_2), the interaction between temperature and volume of ethanol (X_1X_3), and the interaction between time and volume of ethanol (X_2X_3) did not give any significant contribution on Y_1 .

The Lack of Fit F -value of 1.26 implies the Lack of Fit is not significant relative to the pure error. Non-significant lack of fit is good, which indicates that the model is suitable to describe the effect of variables for Y_1 and that the model is adequate for predicting the response. According to Bas and Boyaci (2007), the model will be considered appropriate if the lack of fit value model is not significantly different at the level of specific α . A model will be well fitted to the experimental data if it presents a significant regression and a non-significant lack of fit (Bezerra *et al.*, 2008).

Y_1 was calculated at about 16.38%. The value is comparable to those findings by Anwar *et al.* (2005) of about 15-20% and Pourali *et al.* (2009) of about 10-26% but slightly higher than reported by Mas'ud *et al.* (2017) of about 14.47%. Mas'ud *et al.* (2017) reported the effects of extraction time, temperature, and volume of solvent on mango seed kernel oil above room temperature had a significant effect on oil yield.

3.2.2 γ -Oryzanol of rice bran oil

According to the ANOVA of Y_2 , the F -value model

Table 3. ANOVA for the response variables

Source	df	Oil Yield		γ -oryzanol		TPC	
		SS	p-value	SS	p-value	SS	p-value
Model ^a	9	18.58	6.11E-05	7070.22	0.0001	0.34	5.20E-05
X ₁	1	0.64	0.0445	826.35	0.0029	0.04	0.0013
X ₂	1	2.87	0.0006	979.54	0.0016	0.02	0.0222
X ₃	1	5.11	6.93E-05	894.09	0.0022	0.05	0.0009
X ₁ X ₂	1	0.03	0.6294	2.17	0.8451	0.04	0.0012
X ₁ X ₃	1	0.53	0.0639	14.61	0.6145	0.004	0.1655
X ₂ X ₃	1	0.27	0.1662	582.94	0.0082	0.01	0.0648
X ₁ ²	1	6.32	2.85E-05	1491.02	0.0004	0.01	0.082
X ₂ ²	1	3.52	0.0003	2026.42	0.0001	0.07	0.0001
X ₃ ²	1	0.59	0.0516	969.98	0.0017	0.12	1.90E-05
Residual	10	1.21		540.58		0.02	
Lack of Fit ^b	5	0.68	0.4027	327.086	0.3255	0.01	0.8565
Pure Error	5	0.54		213.49		0.02	
Cor Total	19	19.79		7610.79		0.361207	

^a Significant, ^b Non-significant, X₁: Temperature (°C), X₂: Extraction time(h), X₃: Volume of ethanol (mL)

of 14.53 implies the model is significant. Values of Prob > F of 0.0001 (less than 0.0500) indicate model terms are significant. Variables that have a significant effect on Y₂ are X₁, X₂, X₃, X₂X₃, X₁², X₂², and X₃². Values greater than 0.1000 indicate the model terms are not significant, and that X₁X₂ and X₁X₃ did not give any significant contribution to γ -oryzanol concentration. The Lack of Fit F-value of 1.53 implies the Lack of Fit is not significant relative to the pure error. This indicates that the model is suitable to describe the effect of a parameter observed on γ -oryzanol and that the model is adequate for predicting the response.

γ -oryzanol was extracted at about 783.65 mg. L⁻¹. The value is comparable to those findings by other studies. Arab *et al.* (2011) reported approximately 0.9-2.9% (9.000 - 29.000 mg.L⁻¹), 1.5-2.9% (15.000 - 29.000 mg.L⁻¹) reported by Krishna *et al.* (2001), even up to 3.000 mg.kg⁻¹ reported by Shin *et al.* (1997) and 119.750-281.950 mg.kg⁻¹ reported by Sukanya *et al.* (2017). According to Iqbal *et al.* (2005), the exact composition of γ -oryzanol depends on the rice cultivars. Furthermore, according to Butsat and Siriamornpun (2010), the content of γ -oryzanol in rice is affected by the variety and growing conditions, as the antioxidant component will respond differently to environmental changes.

3.2.3 Total phenolic content for rice bran oil

Based on the ANOVA of Y₃ showed that the F-value model of 17.66, value implies the model is significant. Values of Prob > F of 0.0001 (less than 0.0500) indicate model terms are significant. In this case, X₁, X₂, X₃, X₁X₂, X₂², and X₃². Values greater than 0.1000 indicate the model terms are not significant, therefore X₁X₂ and

X₁X₃ did not give any significant contribution to Y₃. The Lack of Fit F-value of 1.53 implies the Lack of Fit is not significant relative to the pure error, this indicates that the model is suitable to describe the effect of a variable observed for Y₃ and that the model is adequate for predicting. This study obtained a TPC of RBO of 1.02 mg GAE.g⁻¹, this value is in accordance with Bopitiya and Madhujith (2014), they are reported that RBO contains TPC of about 0.8931- 1.2239 mg GAE/mg of bran.

3.3 Interpretation of response surface and contour plots

Based on the ANOVA and the fitted models, the responses surfaces were generated by the model for extraction of Y₁, Y₂, and Y₃ as responses. To gain a better understanding of this study, related to the effect of observed variables on the responses, then the predicted models are presented as the 3D plot (Figures 1-3). Response surface plot is a representation of the surface plot in 3-D space as the plot determining optimum operating conditions reaching maximum from the best-fitted model. These plots are obtained by depicting two variables within the experimental range and keeping the third variable at a constant level. According to Bezerra *et al.* (2008), a two-dimensional representation of a three-dimensional plot can be explained. Thus, if there are three or more variables, the plot visualization is possible only if one or more variables are set at a constant value.

Figure 1 shows a 3D plot corresponding to the effect of X₁ and X₂ on Y₁ at the fixed of X₃. Y₁ had a maximum point, and the effect of X₂ is stronger than the effect of X₁ on increasing Y₁. In the 3D plot curve, the curve at X₂ is more convex than the curve at X₁, meaning that a small change in X₂ has greatly affected the acquisition of

Y_1 compared to the change in X_1 . It is evident in the coefficient variable of X_1 and X_2 , where the coefficient variable of $X_2 > X_1$, and it is evident in the coefficient estimate of the ANOVA, where the coefficient variable in terms of actual factor of X_2 (9.48) is higher than X_1 (1.21). On the effects of X_1 and X_3 on increasing Y_1 at a fixed of X_2 (1.6 h), the coefficient variable in terms of the actual factor of X_1 (1.21) is higher than X_3 (0.61), whereas the effects of X_2 and X_3 on increasing Y_1 at a fixed of X_1 , the coefficient variable in term of the actual factor of X_2 (9.48) is higher than X_3 (0.61).




Figure 1. Response surface 3D plot the effect of X_1 and X_2 on Y_1 at the fixed of X_3

Figure 2 shows a 3D plot corresponding to the effect of X_1 and X_2 on Y_2 at the fixed of X_3 . It can be seen that Y_2 had a maximum point, and the effect of X_2 is stronger than the effect of X_1 on increasing Y_2 . In the 3D plot curve, can be seen that the curve at X_2 is more convex than the curve at X_1 , meaning that a small change in X_2 has greatly affected the acquisition of Y_2 compared to X_1 . It's evident in the coefficient variable of X_1 and X_2 , where the coefficient variable of $X_2 > X_1$, and it is evident in the coefficient estimate of ANOVA, where the coefficient variable in terms of actual factor of X_2 (8.47) is higher than X_1 (7.78). On the effects of X_1 and X_3 on increasing Y_2 at a fixed of X_2 , the effect of X_2 is stronger than the effect of X_1 on increasing Y_2 , the coefficient variable in terms of actual factor of X_1 (7.78) is lower than X_3 (8.09), whereas the effects of X_2 and X_3 on increasing Y_2 at a fixed of X_1 , the effect of X_2 is stronger than the effect of X_3 on increasing Y_2 , the coefficient




Figure 2. Response surface 3D plot the effect of X_1 and X_2 on Y_2 at the fixed of X_3

variable in term of the actual factor of X_2 (8.47) is higher than X_3 (8.09).

Figure 3 shows a 3D plot corresponding to the effect of X_1 and X_2 on Y_3 at the fixed of X_3 . It can be seen that Y_3 had a maximum point, and the effect of X_2 is stronger than the effect of X_1 on increasing Y_3 . A 3D plot curve can be seen that the curve X_2 is more convex than the curve X_1 , meaning that a small change in X_2 has greatly affected the acquisition of Y_3 compared to X_1 , it is evident in the coefficient variable of X_1 and X_2 , where the coefficient variable of $X_2 > X_1$, and it is evident in the coefficient estimate of ANOVA, where the coefficient variable in term of the actual factor of X_2 (0.03) is higher than X_1 (0.06). On the effects of X_1 and X_3 on increasing Y_3 at a fixed of X_2 , can be seen that Y_3 had a maximum point, and the effect of X_1 is lower than the effect of X_3 on increasing Y_3 . At a 3D plot curve, can be seen that curve X_3 is more convex than the curve X_1 , meaning that a small change in X_3 has greatly affected the acquisition of Y_3 compared to X_1 , its evident in the coefficient variable of $X_1 < X_3$, and it is evident in the coefficient estimate of ANOVA, where the coefficient variable in term of the actual factor of X_1 (0.05) is lower than X_3 (0.06), whereas the effects of X_2 and X_3 on increasing Y_3 at a fixed of X_1 , the coefficient variable in term of the actual factor of X_2 (0.03) is lower than X_3 (0.06).




Figure 3. Response surface 3D plot the effect of X_1 and X_2 on Y_3 at the fixed of X_3

3.4 Interpretation of the optimum conditions and verification

According to ANOVA, the optimum of X_1 , X_2 , and X_3 to obtain the maximum of Y_1 , Y_2 , and Y_3 can be achieved at X_1 of 78.7°C, X_2 of 1.6 hrs, and X_3 of 496.5 mL, under the conditions of the extraction process, the predicted values of Y_1 , Y_2 , and Y_3 were 16.4%, 195.49 mg.L⁻¹, and 1,019 mg GAE.g⁻¹, respectively. It can be explained that an increase X_1 of 70°C to 78.7°C, X_2 of 1 h to 1.6 hrs, and X_3 of 350 mL to 496.5 mL at fixed X_1 of 80°C, X_2 of 1.5 hrs, and X_3 of 450 mL cause an increase Y_1 , Y_2 and Y_3 . Further, the addition of X_1 of 78°C up to 90°C, X_2 of 1.6 hrs up to 2 h, and X_3 of 496.5 mL up to

550 mL did not cause an increase in Y_1 , Y_2 and Y_3 .

Loypimai et al. (2020) have noted an increase in extraction temperature caused an increase in the extraction of γ -oryzanol. Previously, Srisaipet and Nuddagu (2014) reported that the amount of γ -oryzanol did not vary significantly at a temperature lower than 120°C, a decrease in γ -oryzanol concentration above 120°C due to the degradation of γ -oryzanol in the oil. According to Nystrom et al. (2007), the good heat stability of RBO has been attributed to its high content of steryl ferulates and tocopherols present in the oil. Related to TPC, Ghafar et al. (2019) reported that the TPC of fruits showed a significant decrease during the heating process at 60, 80, 90, 110, and 130°C. Garas-Salinas et al. (2012) reported that the TPC was detected at 100°C and 5 mins, higher temperatures and longer exposure times reduced extract polyphenol diversity.

For verification models, the laboratory scale has been carried out by conducting triplicate experiments using the recommended/predicted values of X_1 , X_2 and X_3 from the Design Expert 10 software. The result showed that the average of Y_1 , Y_2 , and Y_3 were $16.1 \pm 0.16\%$, $189.12 \pm 1.18 \text{ mg.L}^{-1}$, and $0.91 \pm 0.54 \text{ mg GAE.g}^{-1}$, respectively. The RBO produced in the verification process showed that the average value of Y_1 , Y_2 , and Y_3 was close to the predicted values of the response, although slightly lower than the predicted values of the response. These results indicate that the reflux method of RBO extraction with ethanol as the "green solvent" is feasible for industrial application.

4. Conclusion

The effect of temperature, time, and volume of ethanol as a solvent on the oil yield, γ -oryzanol, and TPC of RBO with apply reflux extraction method have been studied and proven to be able to produce RBO with high oil yield, γ -oryzanol, and TPC. The maximum oil yield = 16.4%, γ -oryzanol = 195.49 mg.L^{-1} , and TPC = $1.02 \text{ mg GAE.g}^{-1}$ were achieved under the optimum conditions of temperature, time, and volume of ethanol were 78.7°C, 1.6 hrs, and 496.5 mL ethanol/100 g rice bran, respectively. The average oil yield, γ -oryzanol, and TPC on verified models were $16.1 \pm 0.66\%$, $189.12 \pm 1.18 \text{ mg.L}^{-1}$, and $0.91 \pm 0.54 \text{ mg GAE.g}^{-1}$, respectively. This result indicates that ethanol is very promising for RBO extraction, as an effort to apply the concept of "green" solvents that are safe for humans and the environment. The optimization of the RBO extraction process has resulted in an extraction model and has been verified on laboratory scale with satisfactory results. The predicted value of oil yield, oryzanol, and TPC close to the value of the verification results have proven that this RBO extraction model is quite valid and feasible to be applied

to the RBO extraction process in the industry.

Conflict of interest

The authors declare no conflict of interest.

Acknowledgements

We would like to thank the Directorate General of Higher Education, Ministry of Education Indonesia, for their research grant No. B/549/PL10/PT.01.05/2021, and the Chemical Engineering Department, Politeknik Negeri Ujung Pandang, Makassar, Indonesia.

References

- Amiri, H., Nabizadeh, R. and Martinez, S.S. (2018). Response surface methodology modeling to improve degradation of Chlorpyrifos in agriculture runoff using TiO₂ solar photocatalytic in a raceway pond reactor. *Ecotoxicology and Environmental Safety*, 147(1), 919-925. <https://doi.org/10.1016/j.ecoenv.2017.09.062>
- Anwar, F., Tabreez, A. and Zahid, M. (2005). Methodical characterization of rice (*Oryza sativa*) bran oil from Pakistan. *Grasas Aceites*, 56(2), 25-134. <https://doi.org/10.3989/gya.2005.v56.i2.120>
- Arab, F., Alemzadeh, I. and Maghsoudi, V. (2011). Determination of antioxidant component and activity of rice bran extract. *Scientia Iranica*, 18(6), 1402-1406. <https://doi.org/10.1016/j.scient.2011.09.014>
- Bas, D. and Boyaci, I.H. (2007). Modelling and optimization I: Usability of response surface methodology. *Journal Food Engineering*, 78(3), 836-845. <https://doi.org/10.1016/j.jfoodeng.2005.11.024>
- Bezerra, M.A., Santelli, R.E., Oliveira, E.P., Villar, L.S. and Escalera, L.A. (2008). Response surface methodology (RSM) as a tool for optimization in analytical chemistry. *Talanta*, 76(5), 965-977. <https://doi.org/10.1016/j.talanta.2008.05.019>
- Bopitiya, D. and Terrence, M. (2014). Antioxidant potential of rice bran oil prepared from red and white rice. *Tropical Agricultural Research*, 26(1), 1-11. <https://doi.org/10.4038/tar.v26i1.8067>
- Brand-Williams, W., Cuvelier, M.E. and Berset, C. (1995). Use of free radical method to evaluate antioxidant activity. *LWT- Food Science and Technology*, 28(1), 25-30. [https://doi.org/10.1016/S0023-6438\(95\)80008-5](https://doi.org/10.1016/S0023-6438(95)80008-5)
- Butsat, S. and Siriamornpun, S. (2010). Antioxidant capacities and phenolic compounds of the husk, bran and endosperm of Thai rice. *Food Chemistry*, 119(2), 606-613. <https://doi.org/10.1016/j.foodchem.2009.07.001>

- Chanioti, S., Siamandoura, P. and Tzia, C. (2016). Evaluation of extracts prepared from olive oil by-products using microwave-assisted enzymatic extraction: effect of encapsulation on the stability of final products. *Waste and Biomass Valorization*, 7 (4), 831-842. <https://doi.org/10.1007/s12649-016-9533-1>
- Do, Q.D., Angkawijaya, A.E., Tran-Nguyen, P.L., Huynh, L.H., Soetaredjo, F.E, Ismadji, S. and Ju, Y.H. (2014). Effect of extraction solvent on total phenol content, total flavonoid content, and antioxidant activity of *Limnophila aromatica*. *Journal of Food and Drug Analysis*, 22(3), 296-302. <https://doi.org/10.1016/j.jfda.2013.11.001>
- Ghafoor, K., Ahmed, I.A.M., Doğu, S., Uslu, N., Fadimu, G.J., Juhaimi, F.A., Babiker, E.E. and Özcan, M.M. (2019). The Effect of Heating Temperature on Total Phenolic Content, Antioxidant Activity, and Phenolic Compounds of Plum and Mahaleb Fruits. *International Journal of Food Engineering*, 15, 11-12. <https://doi.org/10.1515/ijfe-2017-0302>
- Iqbal, S., Bhangar, M.I. and Anwar, F. (2005). Antioxidant properties and components of some commercially available varieties of rice bran in Pakistan. *Food Chemistry*, 93(2), 265-272. <https://doi.org/10.1016/j.foodchem.2004.09.024>
- Krishna, A.G.G., Khatoon, S., Shiela, P.M., Sarmandala, C.V., Indirab, T.N. and Mishrac, A. (2001). Effect of refining of crude rice bran oil on the retention of oryzanol in the refined oil. *Journal of the American Oil Chemists' Society*, 78(2), 127-131. <https://doi.org/10.1007/s11746-001-0232-0>
- Lerma-García, M.J., Herrero-Martínez, J.M., Simó-Alfonso, E.F., Mendonça, C.R.B. and Ramis-Ramos, G. (2009). Composition, industrial processing, and applications of rice bran γ -oryzanol. *Food Chemistry*, 115(2), 389-404. <https://doi.org/10.1016/j.foodchem.2009.01.063>
- Li, L., Li, X. and Yan, C. (2014). Optimization of methyl orange removal from aqueous solution by response surface methodology using spent tea leaves as an adsorbent. *Frontiers of Environmental Science and Engineering*, 8(4), 496-502. <https://doi.org/10.1007/s11783-013-0578-0>
- Loypimai, P., Moongngarm, A., Sittisuanjik, A. and Khamanan, S. (2020). Optimization of tocopherols and γ -oryzanol extraction from rice bran using ultrasound and soybean oil as a green solvent. *Food Research*, 4 (6), 2322-2332. [https://doi.org/10.26656/fr.2017.4\(6\).360](https://doi.org/10.26656/fr.2017.4(6).360)
- Mas'ud, F., Mahendradatta, M., Amran, L. and Zainal, Z. (2017). Optimization of mango seed kernel oil extraction using response surface methodology. *Oilseeds and Fats Crops Lipids*, 24(5), D503. <https://doi.org/10.1051/ocl/2017041>
- Mas'ud, F., Fajar, Banggalino, H., Indriati, S., Todingbua', A., Suhardi and Sayuti, M. (2019). Model development to enhance the solvent extraction of rice bran oil. *Oilseeds and Fats Crops and Lipids*, 26, 16. <https://doi.org/10.1051/ocl/2019009>
- Mas'ud, F., Rifai, A. and Sayuti, M. (2020). Mango seed kernel flour (*Mangifera indica*): nutrient composition and potential as food. *Journal of Nutrition*, 26(1), 101-106. <https://doi.org/10.31246/mjn-2019-0082>
- Mas'ud, F., Indriati, S., Todingbua', A., Rifai, A. and Sayuti, M. (2021). Mango Seed Kernel Oil Extraction with Ethanol: Optimization of Oil Yield and Polyphenol. *Chemical Industry and Chemical Engineering Quarterly*, 27(3), 207-214. <https://doi.org/10.2298/CICEQ200128039M>
- Minjares-Fuentes, R., Femeni, A., Garau, M.C., Meza-Velázquez, J.A., Simala, S. and Rosselló, C. (2014). Ultrasound-assisted extraction of pectin from grape pomace using citric acid: a response surface methodology approach. *Carbohydrate Polymers*, 15 (106), 179-189. <https://doi.org/10.1016/j.carbpol.2014.02.013>
- Mohammed, B.S., Khed, V.C. and Nuruddin, M.F. (2018). Rubbercrete mixture optimization using response surface methodology. *Journal of Clean Production*, 171, 1605-1621. <https://doi.org/10.1016/j.jclepro.2017.10.102>
- Nystrom, L., Achrenius, T., Lampi, A., Moreau, R.A. and Piironen, V. (2007). A comparison of the antioxidant properties of steryl ferulates with tocopherol at high temperatures. *Food Chemistry*, 101(3), 947-954. <https://doi.org/10.1016/j.foodchem.2006.02.046>
- Oliveira, R., Oliveira, V., Aracava, K.K. and Rodrigues, C.E.dC. (2012). Effects of the extraction conditions on the yield and composition of rice bran oil extracted with ethanol - A response surface approach. *Food and Bioprocess Processing*, 90(1), 22-31. <https://doi.org/10.1016/j.fbp.2011.01.004>
- Patel, M. and Naik, S.N. (2004). Gamma-oryzanol from rice bran oil-A review. *Journal of Scientific and Industrial Research*, 63, 569-578.
- Péres, V.F., Maria, J.S. and Melecchi, I.S. (2006). Comparison of soxhlet, ultrasound-assisted and pressurized liquid extraction of terpenes, fatty acids and Vitamin E from *Piper gaudichaudianum* Kunth. *Journal of Chromatography*, 1105(1-2), 115-118.

- <https://doi.org/10.1016/j.chroma.2005.07.113>
- Pourali, O., Feridoun, S.A. and Hiroyuki, Y. (2009). Simultaneous rice bran oil stabilization and extraction using sub-critical water medium. *Journal of Food Engineering*, 95(3), 510-516. <https://doi.org/10.1016/j.jfoodeng.2009.06.014>
- Rodrigues, S., Pinto, G. and Fernandes, F. (2008). Optimization of ultrasound extraction of phenolic compounds from coconut (*Cocos nucifera*) shell powder by response surface methodology. *Ultrasonics Sonochemistry*, 15(1), 95-100. <https://doi.org/10.1016/j.ultsonch.2007.01.006>
- Rodrigues, C.E.C. and Oliveira, R. (2010). Response surface methodology applied to the analysis of rice bran oil extraction process with ethanol. *International Journal of Food Science and Technology*, 45(4), 813-820. <https://doi.org/10.1111/j.1365-2621.2010.02202.x>
- Sani, I. (2014). Soxhlet extraction and physicochemical characterization of *Mangifera indica* L. Seed kernel oil. Research and Reviews. *Journal of Food Dairy Technology*, 2(1), 20-24.
- Shin, T., Godber, J.S., Martin, D.E. and Wells, J.H. (1997). Hydrolytic stability and changes in E vitamers and oryzanol of extruded rice bran during storage. *Journal of Food Science*, 62(4), 704-708. <https://doi.org/10.1111/j.1365-2621.1997.tb15440.x>
- Singh, V., Belova, L., Singh, B. and Sharma, Y.C. (2018). Biodiesel production using a novel heterogeneous catalyst, magnesium zirconate ($Mg_2Zr_5O_{12}$): Process optimization through response surface methodology (RSM). *Energy Conversion and Management*, 174, 198-207. <https://doi.org/10.1016/j.enconman.2018.08.029>
- Srisaipet, A. and Nuddagu, M. (2014). Influence of Temperature on Gamma-Oryzanol Stability of Edible Rice Bran Oil during Heating. *International Journal of Chemical Engineering and Applications*, 5(4), 303-306. <https://doi.org/10.7763/IJCEA.2014.V5.398>
- Sukanya, M., Aikkarach, K., Khongsak, S. and Riantong, S. (2017). Physicochemical and antioxidant properties of rice bran oils produced from colored rice using different extraction methods. *Journal of Oleo Science*, 66(6), 565-572. <https://doi.org/10.5650/jos.ess17014>
- Tan, Y.H., Abdullah, M.O., Hipolito, C.N., Syuhada, N. and Zauzi, A. (2017). Application of RSM and Taguchi methods for optimizing the transesterification of waste cooking oil catalyzed by solid ostrich and chicken-eggshell derived CaO. *Renewable Energy*, 114(Part B), 437-447. <https://doi.org/10.1016/j.renene.2017.07.024>
- Vergara-Salinas, J.R., Perez-Jimenez, J., Torres, J.L., Agosin, E. and Perez-Correa, J.R. (2012). Effects of Temperature and Time on Polyphenolic Content and Antioxidant Activity in the Pressurized Hot Water Extraction of Deodorized Thyme (*Thymus vulgaris*). *Journal of Agricultural and Food Chemistry*, 60(44), 10920-10929. <https://doi.org/10.1021/jf3027759>
- Yolmeş, M., Habibi, N.M.B. and Farhoosh, R. (2014). Optimisation of ultrasound-assisted extraction of natural pigment from annatto seeds by response surface methodology (RSM). *Journal Food Chemistry*, 155, 319-324. <https://doi.org/10.1016/j.foodchem.2014.01.059>

ORIGINALITY REPORT

58%

SIMILARITY INDEX

37%

INTERNET SOURCES

55%

PUBLICATIONS

15%

STUDENT PAPERS

PRIMARY SOURCES

- 1** Fajriyati Mas'ud, Fajar, Herman Bangngalino, Sri Indriati, Abigael Todingbua, Suhardi, Muhammad Sayuti. "Model development to enhance the solvent extraction of rice bran oil", OCL, 2019
Publication **31%**
 - 2** pdfs.semanticscholar.org
Internet Source **5%**
 - 3** 1library.net
Internet Source **2%**
 - 4** Submitted to SDM Universitas Gadjah Mada
Student Paper **2%**
 - 5** Ramon Oliveira, Vinicius Oliveira, Keila Kazue Aracava, Christianne Elisabete da Costa Rodrigues. "Effects of the extraction conditions on the yield and composition of rice bran oil extracted with ethanol—A response surface approach", Food and Bioproducts Processing, 2012
Publication **2%**
-

6	ache.org.rs Internet Source	1 %
7	www.thepharmajournal.com Internet Source	1 %
8	Submitted to Higher Education Commission Pakistan Student Paper	1 %
9	www.tandfonline.com Internet Source	1 %
10	doaj.org Internet Source	1 %
11	Jing-Bo Guo, Ying Fan, Wen-Jie Zhang, Hao Wu, Li-Ming Du, Yin-Xia Chang. "Extraction of gingerols and shogaols from ginger (Zingiber officinale Roscoe) through microwave technique using ionic liquids", Journal of Food Composition and Analysis, 2017 Publication	1 %
12	www.science.gov Internet Source	1 %
13	Submitted to Program Pascasarjana Universitas Negeri Yogyakarta Student Paper	<1 %
14	Soucek, M.D.. "Ternary evaluation of UV-curable seed oil inorganic/organic hybrid	<1 %

coatings using experimental design", Progress in Organic Coatings, 20041220

Publication

15

text-id.123dok.com

Internet Source

<1 %

16

jurnal.poliupg.ac.id

Internet Source

<1 %

17

www.researchgate.net

Internet Source

<1 %

18

Peyman Jalilvand, Ahmad Rahbar-Kelishami, Toraj Mohammadi, Hadi Shayesteh.

"Optimizing of malachite green extraction from aqueous solutions using hydrophilic and hydrophobic nanoparticles", Journal of Molecular Liquids, 2020

Publication

<1 %

19

res.mdpi.com

Internet Source

<1 %

20

www.ars.usda.gov

Internet Source

<1 %

21

iwaponline.com

Internet Source

<1 %

22

Submitted to National Institute of Technology, Rourkela

Student Paper

<1 %

23

Zeng-Xiang Lin, Hong-Man Zhang, Xiao-Jun Ji, Jing-Wen Chen, He Huang. "Hydrolytic Enzyme of Cellulose for Complex Formulation Applied Research", Applied Biochemistry and Biotechnology, 2010

Publication

<1 %

24

www.pgia.pdn.ac.lk

Internet Source

<1 %

25

Kashif Ghafoor, Isam A. Mohamed Ahmed, Süleyman Doğu, Nurhan Uslu et al. "The Effect of Heating Temperature on Total Phenolic Content, Antioxidant Activity, and Phenolic Compounds of Plum and Mahaleb Fruits", International Journal of Food Engineering, 2019

Publication

<1 %

26

Maryam Helmi, Kambiz Tahvildari, Alireza Hemmati, Parviz Aberoomand azar, Aliakbar Safekordi. "Phosphomolybdic acid/graphene oxide as novel green catalyst using for biodiesel production from waste cooking oil via electrolysis method: Optimization using with response surface methodology (RSM)", Fuel, 2020

Publication

<1 %

27

ifst.pericles-prod.literatumonline.com

Internet Source

<1 %

28	ir.lib.seu.ac.lk Internet Source	<1 %
29	Submitted to Universiti Sains Malaysia Student Paper	<1 %
30	ejmcm.com Internet Source	<1 %
31	A. Balasubramanian. "Optimization of process parameters using response surface methodology for the removal of phenol by emulsion liquid membrane", Polish Journal of Chemical Technology, 01/01/2012 Publication	<1 %
32	Submitted to Universitas Jember Student Paper	<1 %
33	www.quickcompany.in Internet Source	<1 %
34	Monica Joshi, Khushwant S. Yadav, Bala Prabhakar. "Quality by Design Approach for Development and Optimization of Rifampicin Loaded Bovine Serum Albumin Nanoparticles and Characterization", Current Drug Delivery, 2021 Publication	<1 %
35	Saousan Chamali, Houcine Bendaoud, Jalloul Bouajila, Severine Camy, Ezzeddine Saadaoui, Jean-Stephane Condoret, Mehrez Romdhane.	<1 %

"Optimization of accelerated solvent extraction of bioactive compounds from Eucalyptus intertexta using response surface methodology and evaluation of its phenolic composition and biological activities", Journal of Applied Research on Medicinal and Aromatic Plants, 2023

Publication

36

academicjournals.org

Internet Source

<1 %

37

www.esanpedia.oar.ubu.ac.th

Internet Source

<1 %

38

A. G. GOPALA KRISHNA. "UNSAPONIFIABLE MATTER AND OXIDATIVE STABILITY OF COMMERCIALY PRODUCED INDIAN RICE BRAN OILS", Journal of Food Lipids, 12/2003

Publication

<1 %

39

Anis Nashuha Md Sani, Noranizan Mohd Adzahan, Mohammad Rashedi Ismail-Fitry. "Valorization of malaysian tropical fruit seeds: A review of their nutrition, bioactivity, processing and food application", Food Bioscience, 2022

Publication

<1 %

40

M.J. Lerma-García, J.M. Herrero-Martínez, E.F. Simó-Alfonso, Carla R.B. Mendonça, G. Ramis-Ramos. "Composition, industrial processing and applications of rice bran γ -oryzanol",

<1 %

Food Chemistry, 2009

Publication

41 acikerisim.ibu.edu.tr <1 %
Internet Source

42 jpmp.org <1 %
Internet Source

43 www.doiserbia.nb.rs <1 %
Internet Source

44 www.foodandnutritionjournal.org <1 %
Internet Source

45 Emmanuel Ohene Afoakwa. "Chocolate Science and Technology", Wiley, 2010 <1 %
Publication

46 Gan, C.Y.. "Optimization of alcohol insoluble polysaccharides (AIPS) extraction from the Parkia speciosa pod using response surface methodology (RSM)", Carbohydrate Polymers, 20100317 <1 %
Publication

47 Renat I. Zhdanov, Anastasia S. Shmyrina, Tatyana V. Zarubina, Andrey L. Mulyukin et al. " Nature of DNA-bound fatty acids in ", FEMS Microbiology Letters, 2006 <1 %
Publication

48 Seng Bee Ng, Ze Yee Pong, Kian Aun Chang, Yun Ping Neo, Lye Yee Chew, Hock Eng Khoo, <1 %

Kin Weng Kong. " Valorisation of mango () kernel as an ingredient of macaron: sensory acceptance and physicochemical properties ", British Food Journal, 2021

Publication

49

Tiraporn Junyusen, Natthaporn Chatchavanthatri, Pansa Liplap, Payungsak Junyusen, Van Man Phan, Siriwan Nawong. "Effects of Extraction Processes on the Oxidative Stability, Bioactive Phytochemicals, and Antioxidant Activity of Crude Rice Bran Oil", Foods, 2022

Publication

<1 %

50

Valentino Bervia Lunardi, Fransiska Gunawan, Felycia Edi Soetaredjo, Shella Permatasari Santoso et al. "Efficient One-Step Conversion of a Low-Grade Vegetable Oil to Biodiesel over a Zinc Carboxylate Metal–Organic Framework", ACS Omega, 2021

Publication

<1 %

51

docksci.com

Internet Source

<1 %

52

dspace.pondiuni.edu.in

Internet Source

<1 %

53

nova.newcastle.edu.au

Internet Source

<1 %

54

psasir.upm.edu.my

Internet Source

<1 %

55

Vergara-Salinas, José Rodrigo, José Cuevas-Valenzuela, and José R. Pérez-Correa.

"Pressurized hot water extraction of polyphenols from plant material",
Biotechnology of Bioactive Compounds, 2015.

Publication

<1 %

56

D. Ranjan. "Parametric Optimization of Selenite and Selenate Biosorption Using Wheat Bran in Batch and Continuous Mode",
Journal of Chemical & Engineering Data,
11/11/2010

Publication

<1 %

57

Fajriyati Mas'ud, Meta Mahendradatta, Amran Laga, Zainal Zainal. "Optimization of mango seed kernel oil extraction using response surface methodology", OCL, 2017

Publication

<1 %

58

J. F. Fu. "Response surface optimization of phosphorus species adsorption onto powdered alum sludge", Journal of Environmental Science and Health Part A,
01/2008

Publication

<1 %

59

Josiane Hilbig, Victor Rodrigues Alves, Carmen Maria Olivera Müller, Gustavo Amadeu Micke

<1 %

et al. "Ultrasonic-assisted extraction combined with sample preparation and analysis using LC-ESI-MS/MS allowed the identification of 24 new phenolic compounds in pecan nut shell [*Carya illinoensis* (Wangenh) C. Koch] extracts", Food Research International, 2018

Publication

Exclude quotes On

Exclude matches < 5 words

Exclude bibliography On