



## EFFECT OF EXTRACTION TIME ON THE BIOACTIVE COMPOUNDS OF BOTTLE GOURD (*LAGENARIA SICERARIA*) USING GAS CHROMATOGRAPHY-MASS SPECTROMETRY

Muhammad Yusuf<sup>1\*</sup>, Sri Indriati<sup>1</sup>, Nur Fitriani Usdyana Attahmid<sup>2</sup>, Rahmawati Saleh<sup>2</sup> and Akhmad Rifai<sup>1</sup>

<sup>1</sup>Department of Chemical Engineering, Politeknik Negeri Ujung Pandang, Makassar City, Indonesia

<sup>2</sup>Department of Agroindustry, Politeknik Pertanian Negeri Pangkep, Pangkep Regency, Indonesia

Traditional medical systems have always been played an important role in meeting global healthcare needs. Meanwhile, bottle gourd (*Lagenaria siceraria*) is a vegetable that contains health-promoting secondary metabolites. Therefore, this study aims to determine the bioactive compounds profiling of Bottle gourd (*Lagenaria siceraria*) fruit extracts in methanol and chloroform using gas chromatography-mass spectrometry (GC-MS) with variations in extraction time of 10, 20, and 30 min using GC-MS RTX-5 capillary column. A total of 91 compounds were tentatively identified, with 55 found in methanol and 41 in chloroform extract. The 1:2 (v/v) ratio using methanol solvent at 30 min was suggested as the most suitable time for maximum extraction. Several peaks with high area percentages were discovered in the methanolic extract containing key chemical constituents such as stearic, oleic, palmitic, and linoleic acid, as well as Cholesta-4,6-dien-3-one, gamma sitosterol, and Phenol, 2,2'-methylene bis. Meanwhile, the corresponding constituents from chloroform extract include Tetracontane, Dotriacontane, Phenol, 2,2'-methylenebis, esters, and aromatic derivatives. Most of the bioactive compounds were detected between 20-30 min time of extraction. Moreover, fatty acids, methyl and ethyl esters, as well as sterols represent 40% of the total extracts and were dominated by oleic, and palmitic acid, gamma-sitosterol along with its ethyl and methyl esters. Therefore, methanol is recommended as the optimal solvent to obtain high content of phytochemical constituents and antioxidants for utilization in pharmacognosy.

**Keywords:** Bottle gourd, Bioactive compounds, GC-MS, Ultra-sonication assisted extraction

### INTRODUCTION

Plants with high antioxidant levels, such as vitamin C, tocopherols, polyphenols and carotenoids, are gaining popularity in the food industry as alternatives to synthetic antioxidants which have limited use due to safety concerns<sup>1</sup>. Meanwhile, synthetic antioxidants have long been used for foods to prevent lipid oxidative rancidity, nutritional loss, off-flavor, quality loss, and discoloration. Aside from extending the shelf life of foods, these compounds also slow the progression of various oxidative stress-related chronic diseases in humans. Furthermore, due to the role in protecting the body from reactive

nitrogen species, crystallization, reactive oxygen species, and free radicals from either normal metabolic processes or external sources, dietary antioxidants play an essential role as nutraceuticals<sup>2-4</sup>. Several mechanisms are presumably involved in this protection, including inhibition of free radical generation, increased scavenging capacity against free radicals, reduced capacity, and metal chelating ability. These reactions are commonly used in antioxidant activity tests. A wide range of activities is determined using antioxidant activity assays with the lipidic system as a substrate<sup>3&5</sup>.

Bottle gourd (*Lagenaria siceraria*) is relatively easy to plant and the planting area is

spread in various parts of the world, ranging from tropical to subtropical climates, as well as highlands to the lowlands. This plant is rich in nutrients containing calcium, iron, vitamin C, polyphenols and saponins which are beneficial for health, therefore, it is taken as daily food. Furthermore, bottle gourd is a common vegetable due to its high choline, phenolics, vitamin B complex, and vitamin C content<sup>6</sup>, while the juice is well-known for its cardioprotective, cardiostimulant, aphrodisiac, and diuretic properties, as well as an antidote to some poisons. Bottle gourd juice is also beneficial for maintaining the body's alkaline reserve due to its less acidic nature<sup>7</sup>. Bottle gourd contains phytochemicals that are beneficial to the body and also produce reactive oxygen species. Meanwhile, the inhibition of reactive oxygen species (ROS) production, direct or indirect scavenging of free radicals, and alteration of intracellular redox potential are all biochemical activities of natural antioxidants. Furthermore, antioxidants, such as carotenoids, flavonoids, polyphenolics, vitamin A, vitamin C, and vitamin E are abundant in vegetables and fruits, preventing free radical damage and lowering the risk of chronic diseases. Therefore, the consumption of dietary antioxidants from these sources potentially prevents cardiovascular diseases, especially atherosclerosis<sup>8</sup>. Meanwhile, the ability of bottle gourd juice to be used as a health drink, is dependent on the extraction and preservation of functional components such as phenolics, carotenoids, and ascorbic acid. Therefore, the processing method selected is essential due to the presence of heat-sensitive components like phenolics, carotenoids, ascorbic acid, and the perishable nature of the product. To date, no attempt has been made to investigate the effects of processing on bottle gourd juice functional components to store and improve efficiency.

Bottle gourd analysis using the ohmic thermal method with variations in temperature and time combined with Gas and Liquid chromatography-mass spectrometry was used to detect volatile and non-volatile phenolics. The ohmically blanched samples exhibited maximum extraction of phenolics and better color of BG juice compared to other samples<sup>9</sup>, while the free radical scavenging activity of *Lagenaria siceraria* fruit ethanolic extract

using the FRAP method was 1.95 mg/ml<sup>3</sup>. In other studies, a combination of the blanching process and sonication extraction to improve the quality of gourd juice bottle showed significant improvements in the total phenolics (TP), carotenoids, total soluble solids (TSS), and physical stability (PS). Other parameters such as titratable acidity (TA), pH, ascorbic acid (AA), browning index (BI), total plate count (TPC), as well as yeast and mold count experienced a significant decrease<sup>10</sup>. The formulations of blended bottle gourd juice, aonla, lemon, and ginger using response surface methodology (RSM) with minimal thermal process showed quality stability against physicochemical, sensory, and microbiology parameters<sup>11</sup>. Moreover, wild bottle gourd optimization using acetone, ethanol, and methanol solvents with Liquid Chromatography-Mass Spectrometry (LC/MS) analysis found the tetracyclic triterpene-cucurbitacin, as well as other pharmaceutically essential compounds<sup>12</sup>. Comparative study of *Lagenaria siceraria* using Soxhlet, microwave-assisted, and ultrasound-assisted extraction, showed that the ultrasonic and microwave assisted extraction methods had an effect on the high levels of polyphenols found in bottle gourd<sup>13</sup>. Another study performed an in-vitro analysis of wild bottle gourd against antioxidant content, antidiabetic, anti-acetylcholine esterase, and anticancer activities using Reversed-Phase-High Performance Liquid Chromatography (RP-HPLC) and FTIR spectroscopy. It was concluded that wild bottle gourd is a rich source of bioactive metabolites<sup>14&15</sup>.

Therefore, this study aims to investigate and characterize the bioactive compounds in the different crude extracts of Bottle gourd (*Lagenaria siceraria*) to determine the physiological, pharmacological, and flavor. Meanwhile, bottle gourd is a medicinal plant and has been attributed to beneficial health effects, but there are only a few studies related to this topic. However, the effect of extraction time with sonication and solvents variations has not been reported. In general, the analysis of bioactive compounds is usually conducted using gas chromatography-mass spectrometry (GC-MS).

## MATERIAL AND METHODS

### Materials

Bottle gourd were collected from Malino district (South Sulawesi province, Indonesia). The plants were taxonomically identified by the Herbarium Bogoriense, Biology Research Center, Indonesian Institute of Sciences (LIPI), Bogor Indonesia (Fig. 1). The chemicals used include analytical grade hexane (emsure 99%), methanol (emsure 99.8%), and chloroform (supelco 99%) supplied by Merck Millipore (Burlington, Massachusetts, United States). Moreover, the instruments used include Shimadzu 2010 GC-MS, Elma Ultrasonic Cleaner S60H, and Buchi Rotary Rotavapor R-300.



**Fig. 1:** Bottle gour

### Preparation of the extract

Bottle gourd was picked and washed with flowing tap water after separating the fruit into epicarp, mesocarp, and seeds. The fresh fruit was then homogenized, for example, the mesocarp was ground separately in an electric mixer grinder. To extract the sample with a ratio of 1:2, 20 ml of bottle gourd juice was mixed with 40 ml hexane (v/v) and transferred to a conical flask which was then immersed in an ultrasonic bath (Elma Ultrasonic) at 40°C for 20 min. Finally, the filtrate was used for sonication extraction using methanol and chloroform solvents.

### Ultra-sonication assisted extraction (UAE)

20 ml filtrate was transferred to a conical flask containing 40 ml solvent methanol or chloroform (1:2 v/v). Furthermore, all the conical flasks were immersed in an ultrasonic bath (Elma Ultrasonic) with a temperature of 40°C, for 10, 20, and 30 min.

**Table 1:** Extraction time of bottle gourd using *Ultra-Sonication Assisted Extraction (UAE)*.

Solvent	Extraction time (min)
Chloroform	10
	20
	30
Methanol	10
	20
	30


### GC-MS determination

This was carried out using Shimadzu 2010 GC-MS and RTX-5 capillary column (30 mm  $\times$  0.25 mm  $\times$  0.25  $\mu$ m) with a split ratio of 40:1 and a temperature of 70°C, heating rate of 10°C  $\text{min}^{-1}$ , up to 300°C, maintained for 5 min with a total analysis time of 25 min. Helium was used as a carrier gas flowing constantly at 1.0 ml/min and the temperature of the inlet was 280°C, pre-column pressure was 80 kPa, and ionization voltage of 70 eV<sup>16</sup>.

## RESULTS AND DISCUSSION

### Identification of bioactive compounds by GC-MS

The phytochemical constituents of bottle gourd were extracted sequentially using two different organic solvents varying in polarity 4.1 (chloroform) and 5.1 (methanol) with a different time extraction (Table 1), while the chemical constituents were analyzed using gas chromatography–mass spectrometry. Fig. 2 illustrates the chromatograms of two crude extracts, with 100 different identified compounds (Tables 2-3), which were then classified into ten chemical groups based on the common name, retention time (Rt), and percent peak area. The chemical groups identified include esters derived from fatty acids, fatty alcohols, fatty acids (FA), amines, aromatic, phenolics, hydrocarbons, terpenes, and sterols, among others. Furthermore, the bioactive compounds were identified using NIST 2.7 and Willey 8 libraries in GC-MS. The chloroform extract contained the fewest compounds (10) after 10 min extraction time, while the highest (25) was identified in the methanol extract after 30 min (Table 2).



**Fig. 2:** Chromatograms of different crude extracts. (A) methanol and chloroform in 10 min time extraction; (B) methanol and chloroform in 20 min time extraction; (C) methanol and chloroform in 30 min time extraction.

### Resource properties of bottle gourd

#### Terpenes

Terpenes were detected using methanol solvents in extraction times of 10 and 30 min as shown in Table 2. Furthermore, Table 3 shows that terpenes were detected in all bottle gourd extracts using chloroform solvents. The total terpenes using both solvents in 10, 20 and 30 min extraction time represented by 2,6,10,14,18,22-tetracosahexaene, and 2,6,10,15,19,23-hexamethyl-, (all-e), were the most dominant and represented 1.55 and 0.47%

of total peak area for methanol extracts as well as 16.27, 4.88, and 5.47% for chloroform extracts. Meanwhile, 2,6,10,14,18,22-tetracosahexaene, and 2,6,10,15,19,23-hexamethyl-, (all-e) have been detected in several plants. These bioactive compounds were identified as strong drugs with biomedical activities to strengthen the body's resistance, resist fatigue, improve human immunity, protect the liver, and were considered substances with great potential in the nutraceutical and pharmaceutical industries in functional and therapeutic applications<sup>17</sup>.

**Table 2:** GC–MS detection of bioactive compounds from bottle gourd using methanol solvent in 10, 20, and 30 min time of extraction.

Peak	R <sub>t</sub>	Area (%)	Bioactive compound
<b>(a) Extraction time 10 min</b>			
<i>Terpenes</i>			
15	33.992	1.55	2,6,10,14,18,22-Tetracosahexaene,2,6,10,15,19,23-hexamethyl-, (all-E)
<i>Esters</i>			
6	16.716	1.63	Propanoic acid, 2-methyl-, 1-(1,1-dimethylethyl)-2-methyl-1,3-propanediyl ester
8	20.918	2.86	Hexadecanoic acid, methyl ester
10	21.843	1.26	Hexadecanoic acid, ethyl ester
11	23.308	0.77	9,12-octadecadienoic acid (z,z)-, methyl ester
12	23.393	0.41	8,11,14-docosatrienoic acid, methyl ester
13	24.291	0.46	Linoleic acid ethyl ester
<i>Fatty alcohols</i>			
7	18.058	0.59	1-hentetracontanol
<i>Fatty acids</i>			
9	21.524	3.18	n-Hexadecanoic acid
<i>Aromatic</i>			
2	5.261	51.86	Ethylbenzene
3	5.433	28.62	P-Xylene
4	5.852	4.18	Benzene, 1,2-dimethyl
<i>Others</i>			
1	5.054	0.73	Cyclotrisiloxane, hexamethyl
5	9.226	0.84	Cyclotrisiloxane, hexamethyl
14	29.924	1.06	Bis(2-ethylhexyl) phthalate
<b>(b) Extraction time 20 min</b>			
<i>Aromatic</i>			
1	5.262	37.59	Ethylbenzene
2	5.333	7.23	Benzene, ethyl-
3	5.433	22.15	P-xylene
4	5.853	1.99	Benzene, 1,2-dimethyl-
<i>Esters</i>			
5	16.712	0.91	Propanoic acid, 2-methyl-, 1-(1,1-dimethylethyl)-2-methyl-1,3-propanediyl ester
7	20.901	2.80	Hexadecanoic acid, methyl ester
8	21.826	1.02	Hexadecanoic acid, ethyl ester
9	23.290	0.85	9,12-octadecadienoic acid (z,z)-, methyl ester
10	24.272	0.63	Linoleic acid ethyl ester
<i>Fatty alcohols</i>			
6	18.018	1.40	1-tetradecanol, acrylate
<i>Phenolics</i>			
11	28.119	2.99	Phenol, 2,2'-methylenebis
<i>Fatty acids</i>			
12	29.913	0.79	1,2-benzenedicarboxylic acid
<i>Sterols</i>			
13	33.314	0.72	Cholesta-4,6-dien-3-one
15	38.965	0.59	Stigmast-5-en-3-ol, oleat
<i>Others</i>			
14	34.021	18.35	Tetrakis (2,3-ditert-butylphenyl)-4,4'-biphenylene diphosphonat

**Table 2:** Continued

<b>(c) Extraction time 30 min</b>			
<i>Aromatic</i>			
1	5.229	20.11	Ethylbenzene
2	5.310	7.24	Benzene, ethyl-
3	5.400	19.44	P-xylene
4	5.823	1.79	Benzene, 1,2-dimethyl-
5	5.942	0.76	Octane, 2,4,6-trimethyl
6	7.794	0.36	Octane, 3,5-dimethyl
<i>Hydrocarbons</i>			
7	9.563	0.37	Undecane
<i>Fatty acids</i>			
8	16.703	0.25	Propanoic acid, 2-methyl-, 1-(1,1-dimethylethyl)-2-methyl-1,3-pro
11	21.437	4.56	n-hexadecanoic acid
17	24.025	2.13	6-octadecenoic acid, (z)
19	24.353	1.35	9,12-octadecadienoic acid (z,z)
23	29.910	0.42	1,2-benzenedicarboxylic acid
<i>Esters</i>			
9	18.339	0.33	Tetradecanoic acid, methyl ester
10	20.883	4.32	Hexadecanoic acid, methyl ester
12	21.811	1.60	Hexadecanoic acid, ethyl ester
13	23.273	1.71	9,12-octadecadienoic acid (z,z)-, methyl ester
14	23.367	1.46	8,11,14-docosatrienoic acid, methyl ester
15	23.446	0.37	9-octadecenoic acid, methyl ester
16	23.742	0.51	Octadecanoic acid, methyl ester
20	24.450	0.37	9-octadecenoic acid (z)-, ethyl ester
21	24.742	0.25	Octadecanoic acid, ethyl ester
18	24.255	1.56	Ethyl (9z,12z)-9,12-Octadecadienoate
<i>Sterols</i>			
22	28.460	26.82	Stigmast-5-En-3-Ol, (3.Beta.,24s)
25	38.976	1.45	Stigmast-5-en-3-ol, oleat
<i>Terpenes</i>			
24	33.986	0.47	2,6,10,14,18,22-tetracosahexaene, 2,6,10,15,19,23-hexamethyl-, (all-e)

**Table 3:** GC-MS detection of bioactive compounds from bottle gourd using chloroform solvent in 10, 20, and 30 min time of extraction.

<b>Peak</b>	<b>R<sub>t</sub></b>	<b>Area(%)</b>	<b>Bioactive compound</b>
<b>(a) Extraction time 10 min</b>			
<i>Esters</i>			
2	20.997	37.36	Hexadecanoic acid, methyl ester
3	21.200	2.16	Beta.-n-acetylneuraminic acid, methyl ester-2-methyl-7,9-methyl-boronate-3,8-di(trimet)
4	21.919	1.57	Hexadecanoic acid, ethyl ester
5	23.466	28.37	9-Octadecenoic acid (Z)-, methyl ester
6	23.831	6.06	Octadecanoic acid, methyl ester
7	29.939	1.51	1,2-benzenedicarboxylic acid, diisooctyl ester
<i>Terpenes</i>			
8	33.995	16.27	2,6,10,14,18,22-Tetracosahexaene, 2,6,10,15,19,23-hexamethyl-, (all-E)- alcohol

**Table 3:** Continued

<i>Aromatic amines</i>			
10	35.283	2.18	1-isopentyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1h-py
<i>Others</i>			
1	6.470	2.53	Ethane, 1,1,2,2-tetrachloro-
9	35.050	1.99	3,3,7,11-Tetramethyltricyclo[5.4.0.0(4,11)]undecan-1-ol
<b>(b) Extraction time 20 min</b>			
<i>Aromatic</i>			
1	5.435	5.58	Ethylbenzene
<i>Fatty acids</i>			
3	16.782	2.28	Propanoic acid, 2-methyl-, 1-(1,1-dimethylethyl)-2-methyl-1,3-pro
7	31.842	1.84	22.alpha.-hydroxy-3,4-secostict-4(23)-en-3-oic acid
<i>Phenolics</i>			
5	28.149	2.78	Phenol, 2,2'-methylenebis[6-(1,1-dimethylethyl)-4-methyl-
<i>Esters</i>			
6	29.935	2.48	1,2-benzenedicarboxylic acid, diisooctyl ester
8	32.025	3.61	Decanoic acid, 8-chloro-, chloromethyl ester
14	34.400	2.86	2,5,9-Trimethyl-12-oxododeca-4,8-dienoic acid, methyl ester
<i>Sterols</i>			
9	32.208	11.56	Stigmast-7-en-3-ol, (3.beta.,5.alpha.,24s)-
10	32.342	3.67	Stigmast-7-en-3-ol, (3.beta.,5.alpha.,24s)-
<i>Terpenes</i>			
12	33.993	48.88	2,6,10,14,18,22-Tetracosahexaene, 2,6,10,15,19,23-hexamethyl-, (all-E)
<i>Hydrocarbons</i>			
13	34.234	4.06	Hexacontane
15	37.556	2.58	Hexacontane
<i>Others</i>			
2	6.472	3.29	Ethane, 1,1,2,2-tetrachloro-
4	18.085	2.49	Spiro(tetrahydrofuryl)2.1'(decalin), 5',5',8'a-trimethyl-
11	33.817	2.04	1-Propanol, 2,3-bis[(3,7,11,15-tetramethylhexadecyl)oxy]-
<b>(c) Extraction time 30 min</b>			
<i>Hydrocarbons</i>			
1	15.419	1.74	Hexadecane, 2,6,10,14-tetramethyl-
17	34.232	6.24	Tetracontane
18	35.765	6.56	Dotriacontane
19	37.570	4.83	Tetracontane
20	39.727	3.54	Dotriacontane
<i>Fatty alcohols</i>			
3	18.103	6.06	1-tetradecanol, acrylate
4	18.404	3.59	1-tridecanol
6	21.583	2.41	1-octadecanol
10	24.777	1.74	1-octadecanethiol
<i>Esters</i>			
2	16.732	8.93	Propanoic acid, 2-methyl-, 1-(1,1-dimethylethyl)-2-methyl-1,3-propanediyl ester
5	20.957	10.31	Hexadecanoic acid, methyl ester
7	21.867	4.60	Hexadecanoic acid, ethyl ester
8	23.413	9.86	9-octadecenoic acid, methyl ester
9	23.782	5.50	Octadecanoic acid, methyl ester

**Table 3:** Continued

11	25.000	2.00	Acetic acid, octadecyl ester
<i>Aromatic</i>			
12	26.813	1.42	1h-purin-6-amine, [(2-fluorophenyl)methyl]-
<i>Phenolics</i>			
13	28.124	11.46	Phenol, 2,2'-methylenebis[6-(1,1-dimethylethyl)-4-methyl-
<i>Terpenes</i>			
16	33.986	5.47	2,6,10,14,18,22-Tetracosahexaene, 2,6,10,15,19,23-hexamethyl-, (all-E)
<i>Others</i>			
14	30.813	2.13	Tetracosamethyl-cyclododecasiloxane
15	32.700	1.61	Tetracosamethyl-cyclododecasiloxane

### Esters

Twenty-two different esters were identified with varying extraction times using methanol solvents (Table 2) and fourteen using chloroform (Table 3). Hexadecanoic acid, methyl and ethyl esters were the most dominant derivative using methanol solvent amounting to 2.86%, 2.80%, and 4.32%, while ester derived from Hexadecanoic acid, ethyl ester represented approximately 1.26%, 1.02%, and 1.60% of the total ester peak area. Meanwhile, the extract with chloroform solvent, identified Hexadecanoic acid, methyl ester (37.36% and 10.31%), 9-Octadecenoic acid (Z)-, methyl ester (28.37% and 9.86%), Propanoic acid, 2-methyl-, 1-(1,1-dimethyl ethyl)-2-methyl-1,3-propanediyl ester (8.93%), Hexadecanoic acid, ethyl ester (1.57% and 4.60%), and Octadecanoic acid, methyl ester (5.50%) as the most dominant derivatives. Other beneficial esters in the extract of bottle gourd, include Octadecanoic acid, methyl ester, and 9-octadecenoic acid, methyl ester. Octadecanoic acid, methyl ester or stearic acid, is suitable for biodiesel production<sup>18</sup>, while 9-octadecenoic acid, methyl ester (oleic acid, methyl ester) reportedly plays an essential role in health care, especially in the treatment of cancer and other diseases<sup>19</sup>. In this case, 9-hexadecenoic acid was used to represent various bottle gourd extracts containing polyunsaturated fatty acids (PUFA/n-7). Besides, PUFAs have been shown to play critical roles in tissue metabolism and cellular, including thermal adaptation, electron and oxygen transport, regulation of membrane fluidity and potentially reduce the risk of coronary heart disease<sup>20</sup>.

### Fatty acids (FA)

Fatty acids are active substrates and allopathic agents with a widely known

antibacterial effect<sup>21</sup>. Saturated fatty acids are synthesized from acetyl coenzyme A by plants and animals as long-term energy storage forms, while saturated fatty acids affect hypercholesterolemia and induce cyclooxygenase-2 expression<sup>22</sup>. As shown in Table 3, the fatty acid composition of bottle gourd, extracted using methanol solvents was represented by five saturated fatty acids namely n-Hexadecanoic Acid (3.18% and 4.56%), 1,2-benzene dicarboxylic acid (0.79% and 0.42%), Propanoic acid, 2-methyl-, 1-(1,1-dimethyl ethyl)-2-methyl-1,3-pro (0.25%), 6-octadecenoic acid, (z) (2.13%), and 9,12-octadecadienoic acid (z,z) (1.35%) (Table 2). Meanwhile, the fatty acid composition identified using chloroform as a solvent showed two saturated fatty acids namely Propanoic acid, 2-methyl-, 1-(1,1-dimethyl ethyl) 2-methyl-1,3-pro (2.28%), and 22.alpha.-hydroxy-3,4-secostict-4(23)-en-3-oic acid (1.84%) (Table 3). n-Hexadecanoic Acid (Palmitic acid) is a saturated long-chain fatty acid with a 16-carbon. Based on the results, palmitic and linoleic acid were the dominant fatty acids bottle gourd extracts. Meat, kernel oil, palm oil, cheese, butter, and milk all contain palmitic acid, this type of fatty acid is reportedly used in pharmaceuticals as an antioxidant, treatment for cancer, and hypercholesterolemic prevention<sup>23</sup>. Meanwhile, Phthalic acid or 1,2-benzenedicarboxylic acid is used in China to clean pollutants and contaminated soils<sup>24</sup>, while Linoleic acid is a source of PUFA, useful in pharmaceutical and medicine for antihistaminic, anticoronary, antieczemic, antiacne, anticancerous, analgesic and ulcerogenic<sup>16&25</sup>.



### Fatty alcohols

Natural fatty alcohols derived from plant or animal lipids are used as detergents, plastics, and in pharmaceuticals<sup>20&26</sup>. The bottle gourd extracted from methanol solvent contains one saturated fatty alcohols namely 1-hentetracontanol (0.59%), while four saturated fatty alcohols namely 1-tetradecanol, acrylate (6.06%), 1-tridecanol (3.59%), 1-octadecanol (2.41%), and 1-octadecanethiol (1.74%) were identified with chloroform (Table 2-3). Saturated fat alcohols are chemical intermediates for surfactants and are widely used in pharmaceutical formulations, agrochemicals, as well as personal, and home care products<sup>20,26</sup>.

### Phenolic compounds

The GC-MS analysis of bottle gourd extracts showed the major phenolics in methanol and chloroform extracts. The percentage of total phenolics varies depending on the extraction solvent, ranging from 2.99% in methanol extract (20 min) to 2.78% and 11.46% in chloroform with an extraction time of 20 and 30 min (Table 2-3). The identified phenolics include Phenol, and 2,2'-methylenebis-6-(1,1-dimethyl ethyl)-4-methyl. Meanwhile, solubility, type of solvent, and polarity in the extraction influence phenolics recovery<sup>27</sup>. Furthermore, the polarity of the solvent is important in increasing solubility<sup>28</sup>. Phenolic compounds have been used pharmacologically as antimicrobial, against neurodegenerative pathologies, and anticarcinogenic. Another study by<sup>29</sup>, reported that the antioxidant capacity of different *Lagenaria siceraria* (bottle gourd) extracts relates directly to the phenolic content. The study identified and isolated six phenolics compounds, including phenolic glycoside (*E*)-4-hydroxymethyl-phenyl-6-*O*-caffeoyl- $\beta$ -d-glucopyranoside which has high antioxidant activity according to the in-vitro analysis. In addition, the fruit of *Lagenaria siceraria* (Molina) is a potentially rich source of natural radical scavengers<sup>3</sup>. Analysis of free radical scavenging activity in ethanol extract showed that the percentage of inhibition was 89.21%.

### Hydrocarbons

The hydrocarbon content in bottle gourd differed between extraction solvents, with Undecane (0.37%) for methanol and

hexacontane (4.06 and 2.58%) for chloroform with an extraction time of 20 min. Meanwhile, for the 30 min, more hydrocarbons were identified namely Hexadecane, 2,6,10,14-tetramethyl (1.74%), Tetracontane (6.24 and 4.83%), and Dotriacontane (6.56 and 3.54%), with the majority being Alkanes (5 hydrocarbons). Tetracontane and Dotriacontane have been detected in *Caralluma retropiciens* (Ehrenb), while *Asclepias Curassavica L* has antimicrobial, antifungal, and antibacterial effects<sup>30&31</sup>.

### Sterols

C29 sterols, also known as phytosterols, are important precursors of vitamin D, while some of the derivatives play a major role in reducing low-density lipoprotein cholesterol in-vivo<sup>32</sup>. Phytosterols in bottle gourd were represented by three different steroids, for an extraction time of 20 min, Cholesta-4,6-dien-3-one (0.72%) and Stigmast-5-en-3-ol, oleat (0.59%) were identified, while more components such as Stigmast-5-en-3-ol, oleat (26.82% and 1.45%) were identified at 30 min. Furthermore, chloroform sterols were identified as Stigmast-7-en-3-ol (11.56 and 3.67%) at total peak area with an extraction time of 20 min (Table 2-3), while stigmast-5-En-3-Ol, (3.Beta.,24s) or Gamma Sitosterol is the most predominant in bottle gourd using methanol solvent. Cholesta-4,6-dien-3-one and gamma sitosterol investigated from Alginate *Glycyrrhiza glabra L* and *Bidens pilosa L* are used in pharmaceuticals as an antihepatotoxic, antiviral, antioxidant, cancer preventive, and hypocholesterolemic<sup>33,34</sup>, while Stigmast-7-en-3-ol investigated from Djulis (*Chenopodium formosanum Koidz.*), has great potential to be developed in the industry as enriched functional foods and nutraceuticals. Based on literature studies, the phytosterols identified by GC-MS are biologically active compounds and have several health benefits, including antioxidant activities and anti-cancer<sup>35</sup>.

### Aromatic, amine and Others

Using methanol extract, two compounds from different chemical groups were identified and are listed in Table 2, namely 1 phthalate derivative (1.06%) and Tetrakis (2,3-ditert-butylphenyl)-4,4'-biphenylene diphosphonat (18.53%), while four compounds were identified from chloroform extract as shown in Table 3,

namely Ethane, 1,1,2,2-tetrachloro (2.53% and 3.29%), Spiro (tetrahydrofuryl)2.1'(decalin), 5',5',8'a-trimethyl (2.49%), and 1-Propanol, 2,3-bis [(3,7,11,15-tetramethylhexadecyl)oxy] (2.04%). In addition, six aromatic compounds were identified in the methanol extract, namely Ethylbenzene (51.86%, 37.59%, and 20.11%), P-xylene (28.62%, 22.15%, and 19.44%), Benzene, 1,2-dimethyl (4.18%, 1.99%, and 1.79%), Benzene, ethyl (4.18%, 7.23%, and 7.24%), Octane, 2,4,6-trimethyl (0.76%), and Octane, 3,5-dimethyl (0.36%) while Ethylbenzene (5.58%) and 1h-purin-6-amine, [(2-fluorophenyl) methyl] (1.42%) were found in the chloroform extract. Also one amines namely 1-isopentyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1h-py (2.18%) was found. Meanwhile, ethylbenzene, p-xylene and benzene, 1,2-dimethyl are aromatic compounds often found in cucurbitaceous plants such as *Lagenaria siceraria* fruits. Tetracosamethylcyclododecasiloxane is commonly used in the cosmetics and fragrance industry<sup>36</sup>, while 3,3,7,11-Tetramethyltricyclo[5.4.0.0(4,11)]undecan-1-ol was detected in *Eucalyptus granlla* wood and is used as pesticide to protect the environment<sup>37</sup>.

### Conclusion

Several valuable compounds were found in the methanol and chloroform extracts of bottle gourd, including fatty acids and alcohols, terpenes, phenolics, hydrocarbons, amines, and esters. The solvent used affects the extracted compounds, particularly fatty acids, esters, amines, phenolics, and sterols, while the presence of antioxidants and sterols (PUFA) in bottle gourd suggests its potential as a source of nutrient in human and animal foods. Furthermore, bottle gourd potential in industrial fragrance and cosmetics was demonstrated by the high concentration of hydrocarbons, esters, and amines. The extraction time of 20-30 min is the optimal range for maximum retention of bioactive compounds. Meanwhile, the identified compounds play an essential role in the development of food functionals and pharmaceutical prospects. However, further studies are needed to identify the various biological activities for the better development of novel pharmaceuticals and food functionals.

### Competing Interests

The authors declare that there are no competing interests.

### Funding

The authors are grateful to Politeknik Negeri Ujung Pandang for the financial support provided through BOPTN in 2020.

### Acknowledgement

The authors are grateful to Politeknik Negeri Ujung Pandang for supporting this study through the BOPTN funding scheme in 2020.

### REFERENCES

1. V. J. Msukwa, C. R. Y. Munthali, B. I. Nyoka and E. Missanjo, "Phenology of sclerocarya birrea (A. rich.) hochst. Provenances", *Emerg Sci J*, 3, 1–13 (2019).
2. D. Bera, D. Lahiri and A. Nag, "Studies on a natural antioxidant for stabilization of edible oil and comparison with synthetic antioxidants", *J Food Eng*, 74, 542-545 (2006).
3. Mayakrishnan, V., Veluswamy, S., Sundaram, K. S., Kannappan, P. and N. Abdullah, "Free radical scavenging potential of *Lagenaria siceraria* (Molina) Standl fruits extract", *Asian Pac J Trop Med*, 6(1), 20–26 (2013).
4. P. H. Kien, Y. Khamphone and G. T. T. Trang, "Study of Effect of Size on Iron Nanoparticle by Molecular Dynamics Simulation", *HighTech Innov J*, 2(3), 158–167 (2021).
5. M. C. Foti, "Antioxidant properties of phenols", *J Pharm Pharmacol*, 59(12), 1673–1685 (2007).
6. S. Bhat, C. S. Saini, M. Kumar and H. K. Sharma, "Effect of Thermal and Alternate Thermal Processing on Bottle Gourd (*Lagenaria siceraria*) Juice", *J Food Process Preserv*, 41, 1–9 (2017).
7. A. Upaganlawar and R. Balaraman, "Cardioprotective effects of *Lagenaria siceraria* fruit juice on isoproterenol-induced myocardial infarction in wistar rats: A biochemical and histoarchitecture study", *J Young Pharm*, 3(4), 297–303 (2011).
8. B. Uttara, A. Singh, P. Zamboni and

- R.Mahajan, "Oxidative Stress and Neurodegenerative Diseases: A Review of Upstream and Downstream Antioxidant Therapeutic Options", *Curr Neuropharmacol*, **7**, 65–74 (2009).
9. S. Bhat, C. S. Saini and H. K. Sharma, "Changes in total phenolic content and color of bottle gourd (*Lagenaria siceraria*) juice upon conventional and ohmic blanching", *Food Sci Biotechnol*, **26**, 29–36 (2017).
  10. S. Bhat and H. K. Sharma, "Combined effect of blanching and sonication on quality parameters of bottle gourd (*Lagenaria siceraria*) juice", *Ultrason Sonochem*, **33**, 182–189 (2016).
  11. R. R. Gajera and D. C. Joshi, "Development and quality evaluation of bottle gourd, *lagenaria siceraria* (Mol.) standl. based blend juice", *Indian J Nat Prod Resour*, **6**, 194–199 (2015).
  12. U. A. Attar and S. G. Ghane, "Optimized extraction of anti-cancer compound – cucurbitacin I and LC–MS identification of major metabolites from wild Bottle gourd (*Lagenaria siceraria* (Molina) Standl.)", *South African J Bot*, **119**, 181–187 (2018).
  13. M. Abbas, D. Ahmed, M. T. Qamar, S. Ihsan and Z. I. Noor, "Optimization of ultrasound-assisted, microwave-assisted and Soxhlet extraction of bioactive compounds from *Lagenaria siceraria*: A comparative analysis", *Bioresour Technol Reports*, **15**, 100746 (2021).
  14. U. A. Attar and S. G. Ghane, "In-vitro antioxidant, antidiabetic, antiacetylcholine esterase, anticancer activities and RP-HPLC analysis of phenolics from the wild bottle gourd (*Lagenaria siceraria* (Molina) Standl.)", *South African J Bot*, **125**, 360–370 (2019).
  15. A. Dutta, N. Roy, K. Das, *et al.*, "Synthesis and characterization of host guest inclusion complexes of cyclodextrin molecules with theophylline by diverse methodologies", *Emerg Sci J*, **4**(1), 52–72 (2020).
  16. M. Yusuf, N. F. Atthamid, S. Indriat, R. Saleh, M. Latife and A. Rifai, "Optimization ultrasonic assisted extraction (UAE) of bioactive compound and antibacterial potential from sea urchin (*diadema setosum*)", *Curr Res Nutr Food Sci*, **8**(2), 556–569 (2020).
  17. W. Peng, *et al.*, "Characteristics of antibacterial molecular activities in poplar wood extractives", *Saudi J Biol Sci*, **24**(2), 399–404 (2017).
  18. A. L. V. Cubas, M. M. Machado, C. R. S. C. Pinto, E. H. S. Moecke and A. R. A. Dutra, "Biodiesel production using fatty acids from food industry waste using corona discharge plasma technology", *Waste Manag*, **47**(ptA), 149–154 (2016).
  19. C. A. Ukwubile, A. Ahmed, U. A. Katsayal, J. Ya'u, and S. Mejida, "GC–MS analysis of bioactive compounds from *Melastomastrum capitatum* (Vahl) Fern. leaf methanol extract: An anticancer plant", *Sci African*, **3**, e00059 (2019).
  20. E. I. Abdel-Aal, A. M. Haroon and J. Mofeed, "Successive solvent extraction and GC-MS analysis for the evaluation of the phytochemical constituents of the filamentous green alga *Spirogyra longata*", *Egypt J Aquat Res*, **41**(3), 233–246 (2015).
  21. L. J. McGaw, A. K. Jäger and J. Van Staden, "Antibacterial effects of fatty acids and related compounds from plants", *South African J Bot*, **68**, 417–423 (2002).
  22. J. Y. Lee, K. H. Sohn, S. H. Rhee and D. Hwang, "Saturated Fatty Acids, but Not Unsaturated Fatty Acids, Induce the Expression of Cyclooxygenase-2 Mediated through Toll-like Receptor 4", *J Biol Chem*, **276**(20), 16683–16689 (2001).
  23. M. Ahmad, W. N. Baba, A. Gani, *et al.*, "Effect of extraction time on antioxidants and bioactive volatile components of green tea (*Camellia sinensis*), using GC/MS", *Cogent Food Agric*, **1**, 1–11 (2015).
  24. T. Ma, *et al.*, "A new procedure combining GC-MS with accelerated solvent extraction for the analysis of phthalic acid esters in contaminated soils", *Front Environ Sci Eng China*, **7**, 31–42 (2013).
  25. E. Vadivel and S. B. Gopalakrishnan, "GC-MS analysis of some bioactive constituents of *Mussaenda frondosa* Linn", *Int J Pharma Bio Sci*, **2**(1), 313–320 (2011).

26. P. Roose, K. Eller, E. Henkes, R. Rossbacher and H. Höke, "Amines, Aliphatic in Ullmann's Encyclopedia of Industrial Chemistry", *Ullmann's Encyclopedia of Industrial Chemistry*, (2015).
27. M. Dent, V. Dragovi-Uzelac, M. Peni *et al.*, "The effect of extraction solvents, temperature and time on the composition and mass fraction of polyphenols in dalmatian wild sage (*Salvia officinalis* L.) extracts", *Food Technol Biotechnol*, 51(1), 84-91 (2013).
28. M. Naczka and F. Shahidi, "Phenolics in cereals, fruits and vegetables: Occurrence, extraction and analysis", *J Pharm Biomed Anal*, 41(5), 1523-1542 (2006).
29. R. Mohan, R. Birari, A. Karmase, S. Jagtap and K. K. Bhutani, "Antioxidant activity of a new phenolic glycoside from *Lagenaria siceraria* Stand", *Fruits Food Chem*, 132(1), 244-251 (2012).
30. S. S. Alqahtani, H. A. Makeen, S. J. Menachery and S. S. Moni, "Documentation of bioactive principles of the flower from *Caralluma retrospiciens* (Ehrenb) and in vitro antibacterial activity – Part B", *Arab J Chem*, 13(10), 7370-7377 (2020).
31. S. Bihana, A. Dhiman, G. Singh and S. Satija, "Gas chromatography-mass spectroscopy analysis of bioactive compounds in the whole plant parts of ethanolic extract of *Asclepias Curassavica* L", *Int J Green Pharm*, 12(2), 107-114 (2018).
32. M. Francavilla, P. Trotta and R. Luque, "Phytosterols from *Dunaliella tertiolecta* and *Dunaliella salina*: A potentially novel industrial application", *Bioresour Technol*, 101(11), 4144-4150 (2010).
33. R. Akhtar, and A. Shahzad, "Alginate encapsulation in *Glycyrrhiza glabra* L. with phyto-chemical profiling of root extracts of in vitro converted plants using GC-MS analysis", *Asian Pac J Trop Biomed*, 7(10), 855-861 (2017).
34. Y. Shen, Z. Sun, P. Shi, *et al.*, "Anticancer effect of petroleum ether extract from *Bidens pilosa* L and its constituent's analysis by GC-MS", *J Ethnopharmacol*, 217, 126-133 (2018).
35. C. Y. Huang, Y. L. Chu, K. Sridhar and P. J. Tsai, "Analysis and determination of phytosterols and triterpenes in different inbred lines of *Djulis* (*Chenopodium formosanum* Koidz.) hull: A potential source of novel bioactive ingredients", *Food Chem*, 297(1), 124948 (2019).
36. N. Fadle, "Abdalbasit Adam Mariod, Hiba Abdel Rahman Ali and Alfatih Ahmed. TLC and GC-MS analysis of fermented wood 'Nikhra' petroleum ether fraction of *Combretaceae* spp. *Combretum hartmannianum* and *Terminalia laxiflora*", *Eurasian J For Sci*, 6(3), 1-7 (2018).
37. S. Ge, W. Peng, D. Li, *et al.*, "Study on antibacterial molecular drugs in *Eucalyptus granilla* wood extractives by GC-MS", *Pak J Pharm Sci*, 28(4 Suppl), 1445-1448 (2015).
38. M. Yusuf, U.A. Fitriani Nur, L. Mahyati, and M. Imran, "Phytochemical and antibacterial properties of sea cucumber (*Muelleria lecanora*) from Barrang Lompo Islands, Makassar South Sulawesi", *Food Res*, 4(6), 1885-1895 (2020).
39. A. H. Shobier, S. A. Abdel Ghani, and K. M. Barakat, "GC/MS spectroscopic approach and antifungal potential of bioactive extracts produced by marine macroalgae", *Egypt J Aquat Res*, 42(3), 289-299 (2016).
40. E. Beyzi, K. Karaman, A. Gunes and S. Buyukkilic Beyzi, "Change in some biochemical and bioactive properties and essential oil composition of coriander seed (*Coriandrum sativum* L.) varieties from Turkey", *Ind Crops Prod*, 109, 74-78 (2017).



## نشرة العلوم الصيدلانية جامعة أسيوط



### تأثير وقت الاستخلاص على المركبات الحيوية من القرع (لاجناريا سيكراريا) باستخدام كروماتوجرافيا الغاز - مقياس الطيف الكتلي

محمد يوسف\*<sup>١</sup> - سري أندرياتي<sup>١</sup> - نور فطراني أسديانا أتاحميد<sup>٢</sup> - رحمواتي صالح<sup>٢</sup> -  
أحمد رفاعي<sup>١</sup>

<sup>١</sup> قسم الهندسة الكيميائية، كلية الفنون التطبيقية، مدينة ماسار، إندونيسيا

<sup>٢</sup> قسم الصناعات الزراعية، كلية الفنون التطبيقية الزراعية، ولاية بانجكيب، ريجنسي بانجكيب، إندونيسيا

لطالما لعبت الأنظمة الطبية التقليدية دورًا مهمًا في تلبية احتياجات الرعاية الصحية العالمية. وفي الوقت نفسه، فإن القرع الزجاجي هو نبات يحتوي على نواتج الأيض الثانوية المعززة للصحة. لذلك، تهدف هذه الدراسة إلى تحديد خصائص المركبات النشطة بيولوجيًا لمستخلصات فاكهة القرع الزجاجي في الميثانول والكلوروفورم باستخدام كروماتوجرافيا الغاز - مقياس الطيف الكتلي للغاز (GC-MS) مع اختلافات في وقت الاستخلاص تبلغ ١٠ و ٢٠ و ٣٠ دقيقة باستخدام GC - عمود شعري RTX-5 MS. تم تحديد إجمالي ٩١ مركبًا مبدئيًا، مع ٥٥ منها في الميثانول و ٤١ في مستخلص الكلوروفورم. تم اقتراح نسبة ١:٢ (حجم / حجم) باستخدام مذيب الميثانول عند ٣٠ دقيقة على أنها أنسب وقت للاستخلاص الأقصى. تم اكتشاف العديد من القمم ذات النسب المئوية العالية في المستخلص الميثانولي الذي يحتوي على مكونات كيميائية رئيسية مثل حامض دهني، أوليك، بالميتيك، ولينوليك، بالإضافة إلى كوليستا٤-٥-دين-٣-اون وجاما سيتوستيرول و فينول ٢، ٢-ميثيلين مكرر، وكذلك تشتمل المكونات من مستخلص الكلوروفورم على تيتراكونتان و دوتراكونتان و دوترياكوتان و فينول ٢، ٢-ميثيلين مكرر والإسترات والمشتقات العطرية. تم الكشف عن معظم المركبات النشطة بيولوجيا بين ٢٠-٣٠ دقيقة من وقت الاستخلاص. علاوة على ذلك، تمثل الأحماض الدهنية، وإسترات الميثيل والإيثيل، وكذلك الستيرويدات ٤٠٪ من إجمالي المستخلصات وتغلب عليها حمض الأوليك والبالميتيك، جاما سيتوستيرول مع إسترات الإيثيل والميثيل. لذلك، يوصى باستخدام الميثانول باعتباره المذيب الأمثل للحصول على نسبة عالية من المكونات الكيميائية النباتية ومضادات الأكسدة لاستخدامها في كعقاقير.