

Biodegradation Of Lignin From Corn Cob By Using A Mixture Of Phanerochaete Chrysosporium, Lentinus Edodes And Pleurotus Ostreatus

by Mahyati Mahyati

Submission date: 27-Jul-2023 08:18PM (UTC-0400)

Submission ID: 2137774482

File name: chaete-Chrysosporium-Lentinus-Edodes-And-Pleurotus-Ostreatus.pdf (455.97K)

Word count: 1955

Character count: 10525

Biodegradation Of Lignin From Corn Cob By Using A Mixture Of Phanerochaete Chrysosporium, Lentinus Edodes And Pleurotus Ostreatus

Mahyati, Abdul Rauf Patong, Muh. Nasir Djide, Dan Paulina Taba

Abstract: Corn cob is agricultural waste containing approximately 76% holocellulose (cellulose and hemicellulose). Cellulose can be converted into bioethanol using a method of SSF. Degradation of lignin from corn cob can be conducted by acid, basic and enzymatic methods. The method applied to the process of enzymatic pretreatment was conducted by using a mixture of white rot fungi (Phanerochaete chrysosporium, Lentinus edodes and Pleurotus ostreatus) to degrade lignin which is environmentally friendly. Results showed that maximum lignin biodegradation, i.e. 96.88%, was achieved after incubation for 30 days by the mixture of white rot fungi

Keyword: biodegradation, lignin, white rot fungi and corn cob

INTRODUCTION

Corn cobs are found as many as 30 b/b% of corn (Koswara, 1991). It is used as an ideal source of lignocellulose which can be found easily. The production of corns in South Sulawesi achieved 424 thousands hectares with the productivity of 4.55 ton per hectare or 1.32 million per year (BPS,2012). The use of corn cobs requires pretreatment (Saha,2003), that can be done by acid, base or enzymatic methods (Isroi, 2011; Alvira et al. 2010; Hendriks and Zeeman 2009; Taherzadeh and Karimi 2008; Hu et al. 2008; Mosier 2005;). An enzymatic pretreatment has several advantages, i.e. enzyme works specifically, therefore sugar produced from hydrolysis will not degraded, the process can be conducted at low temperature and neutral medium, higher product will be obtained, and the maintenance cost will be low because the corrosive materials are not used (Taherzadeh dan Karimi, 2007). Generally, enzymes, which are secondary metabolic from white rot fungi, such as lignin peroxidase (LiP), manganese peroxidase (MnP), laccase (Lac) dan versatile peroxidase (VP). The enzymes play an important role in degradation process of lignin and can oxidize phenolic compounds found in lignocellulose (Van der Merwe, 2002; Higuchi 2004; Wong 2009).

Therefore, the use white rot fungi can be an alternative solution for degrading lignin. Each species of white rote fungi has specific ability in degrading lignin from corn cob. Degradation of lignin was affected by incubation time of fungi and variation of the mixture of white rot fungi (Zeng et al. 2010; Taccari et al. 2009). *P. chrysosporium* is a white rot fungus that can produce LiP and MnP, but it cannot produce Lac enzyme (Howard, et al, 2003). Lac enzyme is produced by *L. edodes* dan VP is obtained from *P. ostreatus* (Dashtban, et al, 2010). Enzyme produced by *P. ostreatus* is MnP (Sarkar dkk. 1997). Some researchers have reported degradation of lignin by using just one white rot fungus. This research used three kinds of white rot fungi (*P. chrysosporium*, *L. edodes*, and *P. ostreatus*) for degrading lignin from corn cobs..

MATERIALS AND METHODS

Corn cobs were obtained from waste of corn plants grown by corn farmers from several districts in South Sulawesi Province. Corn cobs were crushed and the corn cob powder obtained were sieved and the powder passed by the 45 meshed filter was collected and dried until the water content of below 10%.

Preparation of delignification media from corn cobs

Corn cob powder (10 g) was put into a 250 mL Erlenmeyer, 0.01 g of glucose and 5 mL of 0.004% Tween solution were added into the Erlenmeyer, mixed and then sterilized at 121 °C for 15 min in an autoclave. A 5 mL medium of fungi inoculum was put into the sterile mixture, covered with cotton and aluminum foil and incubated at room temperature for 30 days was analyzed every 5 days. The same treatment was conducted for cultures of *P. chrysosporium*, *L. edodes* dan *P. ostreatus*.

Analysis of lignin and cellulose

Analysis of lignin and cellulose followed the method of Chesson (Datta, 1981). A mixture containing 1 g of dried sample (a) and 150 mL of aquadest was heated in a water bath at a temperature of 90 - 100 °C for 1 h. The mixture was filtered and the residue was washed with hot water (300 mL). The residue was dried in the oven until the weight was constant (b). The residue was mixed with 150 mL of 1 N H₂SO₄ and heated in the water bath at 90 - 100 °C for 1 h. The mixture was filtered and washed with 300 mL of

- Mahyati, Abdul Rauf Patong, Muh. Nasir Djide, Dan Paulina Taba
- Department Of Chemical Engineering, State Polytechnic of Ujung Pandang, Indonesia
- Department Of Chemistry, The Faculty of Mathematics and Natural Sciences, Hasanuddin University, Indonesia
- Department of Pharmacy, The Faculty of Pharmacy, Hasanuddin University, Indonesia
- Email; mahyati@yahoo.com
- Telephone +62 085298353527, Fax +62 (0411) 586043

aquades and then the residue was dried (c). The dried residue was soaked with 10 mL of 72% H₂SO₄ at room temperature for 4. After that, 150 mL of 1 N H₂SO₄ was added into the mixture and refluxed in the water bath for 1 h. The solid was washed with 400 mL of aquades, heated in the oven at 105 °C and weighed until the constant weight (d). Finally the solid was heated until become ash and weighed (e). The percentage of cellulose and lignin was calculated as follows:

$$\% \text{ cellulose} = \frac{c-d}{a} \times 100\% \dots (1)$$

$$\% \text{ lignin} = \frac{d-e}{a} \times 100\% \dots (2)$$

where: a = sample weight (g), c = the residue weight (g) at the third weighing, d = the residue weight (g) at the fourth weighing, e = the weight of ash (g).

RESULTS AND DISCUSSION

The reduced of cellulose and lignin content as a function of the incubation time using a mixture of 3 species of the white rot fungi was given in Figure 1. The maximum result of corn

cobs delignification was obtained with the small amount of cellulose reduced because in the presence of the white rot fungi, different enzymes were produced. The smallest lignin content (0.5 %) for the mixture containing 3 species of fungi was obtained at the incubation time of 30 min at the day-30. At the same condition, the cellulose content was 32.92 %. The maximum lignin content reduced was 96.88 %. Therefore at the day-30, the percentage of lignin reduced was 96,88 % whereas the percentage of cellulose reduced was 17,70%. This occurs because with the presence of the white rot fungi, enzymes of oxalate oxidase and oxalate dehydrogenase were produced. The enzymes oxidized oxalic acid and therefore, the cellulose will be degraded. The mixture of 3 species of fungi can produce enzymes optimally that worked together to degrade lignin in corn cobs maximally with the smallest percentage of cellulose reduced. This result showed the maximum initial treatment because lignin cannot be removed completely (100%) due to the presence of bonds between lignin and cellulose that has not been known completely (Steffen, 2003). According to Achmadi (1990), more than 2/3 part of phenyl propane in lignin was related via ether bonds and the rest was related through carbon-carbon bonds.




Figure 1.

The reduced percentage of cellulose and lignin as a function of the incubation time using a mixture of 3 species of the white rot fungi (*P. chrysosporium*, *L. edodes* and *P. ostreatus*)

At the delignification process using inoculum consisting 3 species of white rot fungi, several enzymes can be produced that worked together to degrade lignin. The enzymes working in the ligninolysis of corn cobs was LIP acting as the main catalyst for degrading non-phenolic units, MnP oxidizing Mn²⁺ to Mn³⁺ to break the phenolic unit of lignin. The other enzyme work in the process was Lac (oxidizing enzyme) working through a dimethylation process changing methoxy groups to methanol. The radical

compounds spontaneously or step by step will remove the intermolecular bonds and some of them will destroy the aromatic rings. Therefore, the bond between lignin and cellulose degraded. Figure 2 shows the presence of Lac produced by *L. edodes* that can oxidize phenolic groups to quinon (Arora and Sandhu, 1985) and produce brown or black colour at corn cobs after incubation with the white rot fungi at the day-30.



a). before delignification

b). After delignification

Figure 2.

The product of corn cob delignification using mixed inoculum of the white rot fungi (*P. chrysosporium*, *L. edodes* and *P. ostreatus*)

CONCLUSION

Degradation of lignin using white rot fungi (*P. chrysosporium*, *L. edodes* and *P. ostreatus*) achieved 96.88% and the degradation present age of cellulose was 17,70%.

REFERENCES

- [1]. Koswara, J. Budidaya Jagung, Institut Pertanian Bogor. Bogor (1991)
- [2]. Badan Pusat Statistik Provinsi Sulawesi Selatan. Produksi Padi & Jagung, Makassar (2012)
- [3]. Saha, B. C, Hemicellulose Bioconversion, *J. Ind. Microbio. Biotechnol.*, 30, 279-291 (2003)
- [4]. Isroi, Ria Millati, Siti Syamsiah, Claes Niklasson, Muhammad Nur Cahyanto, Knut Lundquist, and Mohammad J. Taherzadeh, Biological pretreatment of lignocelluloses with white-rot fungi and its applications: A review, *Bioresources* 6(4), Nov. 1 (2011)
- [5]. Alvira, P., Tomás-Pejó, E., Ballesteros, M., and Negro, M. J. Pretreatment for an efficient bioethanol production process based on enzymatic hydrolysis: A review, *Bioresource Technol.* 101, 4851-4861. (2010)
- [6]. Hendriks, A. T. W. M., and Zeeman, G. Pretreatments to enhance the digestibility of lignocellulosic biomass," *Bioresource Technol.* 100, 10-18 (2009)
- [7]. Taherzadeh, M. J., and Karimi, K. "Pretreatment of lignocellulosic waste to improve ethanol and biogas production," *Int. J. Mol. Sci.* 9, 1621-1651 (2008)
- [8]. Hu, G., Heitmann, J. A., and Rojas, O. J. Feedstock pretreatment strategies for producing ethanol from wood, bark and forrest residues, *Bioresources* 3, 270-294 (2008)
- [9]. Mosier, N., Wyman, C., Dale, B., Elander, R., Lee, Y. Y., Holtzapple, M., and Ladisch, M. Features of promising technologies for pretreatment of lignocellulosic biomass. *Bioresource Technol.* 96, 673-686 (2005)
- [10]. Taherzadeh MJ, Karimi K.. Acid-based hydrolysis processes for ethanol from lignocellulosic materials: a review. *Bioresources* 2 (3): 472-499 (2007)
- [11]. Higuchi, T.. Microbial degradation of lignin: Role of lignin peroxidase, manganese peroxidase, and laccase, *Proc. Jpn. Acad., Ser.B* 80, 204-211 (2004)
- [12]. Wong, D. Structure and action mechanism of ligninolytic enzymes, *App. Biochem. and Biotech.* 157, 174-209 (2009).
- [13]. Van der Merwe, J.J., Production of Laccase by The White-Rot Fungus *Pycnoporus sanguineus*, Master Thesis, University of the Free State, Bloemfontein. (2002)
- [14]. Howard RL., E. Abotsi, EL Jansen van Rensburg, S Howard. Lignocellulose biotechnology: issues of bioconversion and enzyme production. *African Biotechnol.* 2 (12): 602-619 (2003)
- [15]. Dashtban, M., Schraft, H., Syed A. A., Qin W. S., Fungal Biodegradation And Enzymatic Modification Of Lignin, Review (2010)
- [16]. Zeng, G., Huang, D., Huang, G., Hu, T., Jiang, X., Feng, C., Chen, Y., Tang, L., and Liu, H. Composting of lead-contaminated solid waste with inocula of white-rot fungus, *Bioresource Technol.* 98, 320-326 (2007).

- [17]. Taccari, M., Stringini, M., Comitini, F., and Ciani, M. Effect of *P. chrysosporium* inoculation during maturation of co-composted agricultural wastes mixed with olive mill wastewater, *Waste Manag.* **29**, 1615-1621 (2009)
- [18]. Datta, R., Acidogenic Fermentation Of Lignocellulose-Acid Yield And Conversion Of Components, *Biotech. and Bioeng.* **23** (9), 2167-2170 (1981)
- [19]. Steffen, K. T. Degradation Of Recalcitrant Biopolymers And Polycyclic Aromatic Hydrocarbons By Litter-Decomposing Basidiomycetous Fungi, Dissertation, Finland, Dept. of App. Chem. and Micr. Viikki Biocenter, University of Helsinki (2003)
- [20]. Achmadi, S.S. Diktat Kimia Kayu. Bogor: Pusat, Antar Universitas, Institut Pertanian Bogor (1990)
- [21]. Arora, D. S., Sandhu. D. K., Laccase production and wood degradation by a white-rot fungus *Daedalea flavida*, *Enzyme Microb. Technol.* **7**, 405-408 (1985)

Biodegradation Of Lignin From Corn Cob By Using A Mixture Of Phanerochaete Chrysosporium, Lentinus Edodes And Pleurotus Ostreatus

ORIGINALITY REPORT

7%

SIMILARITY INDEX

3%

INTERNET SOURCES

6%

PUBLICATIONS

2%

STUDENT PAPERS

PRIMARY SOURCES

1	Submitted to Syiah Kuala University Student Paper	1%
2	silopub Internet Source	1%
3	patents.google.com Internet Source	1%
4	Graciela M.L. Ruiz-Aguilar, José M. Fernández-Sánchez, Refugio Rodríguez-Vázquez, Héctor Poggi-Varaldo. "Degradation by white-rot fungi of high concentrations of PCB extracted from a contaminated soil", Advances in Environmental Research, 2002 Publication	1%
5	Pratima Bajpai. "Biological Bleaching of Chemical Pulps", Critical Reviews in Biotechnology, 1/1/2004 Publication	1%
6	Submitted to Universitas Jember Student Paper	1%

7	Balat, M.. "Progress in bioethanol processing", Progress in Energy and Combustion Science, 200810 Publication	1 %
8	Efri Mardawati, R Andoyo, K A Syukra, MTAP Kresnowati, Y Bindar. "Production of xylitol from corn cob hydrolysate through acid and enzymatic hydrolysis by yeast", IOP Conference Series: Earth and Environmental Science, 2018 Publication	1 %
9	M. Taccari, M. Stringini, F. Comitini, M. Ciani. "Effect of Phanerochaete chrysosporium inoculation during maturation of co- composted agricultural wastes mixed with olive mill wastewater", Waste Management, 2009 Publication	1 %
10	Shen, Xiao-Jun, Bing Wang, Pan-Li Huang, Jia- Long Wen, and Run-Cang Sun. "Effects of aluminum chloride-catalyzed hydrothermal pretreatment on the structural characteristics of lignin and enzymatic hydrolysis", Bioresource Technology, 2016. Publication	1 %
11	docksci.com Internet Source	1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

Biodegradation Of Lignin From Corn Cob By Using A Mixture Of Phanerochaete Chrysosporium, Lentinus Edodes And Pleurotus Ostreatus

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4
